

Pistol®



MAPP 18854

A suspension concentrate containing 250g/L glyphosate and 40g/L diflufenican.

A non-selective herbicide for control of annual and perennial weeds on natural surfaces not intended to bear vegetation, permeable surfaces overlaying soil and hard surfaces (railway ballast only).

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces, or applying the product.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

WASH HANDS AND EXPOSED SKIN before meals and after work.

Environmental protection

Do not contaminate water with the product.

Do not clean application equipment near surface water.

Avoid contamination via drains from farmyards and roads.

Extreme care must be taken to avoid spray drift onto non-crop plants outside the target area.

Hand-held equipment

Since there is a risk to aquatic life from use, direct spray must not be allowed to fall within 2 m of the top of the bank of any static or flowing waterbody or the top of a ditch which is dry at the time of application. Spray must be aimed away from water.

Vehicle-mounted equipment (boom sprayer)

To protect aquatic organisms, respect an unsprayed buffer zone to surface water bodies of 6 m.

HORIZONTAL BOOM SPRAYERS MUST BE FITTED WITH THREE STAR DRIFT REDUCTION TECHNOLOGY.

Low drift spraying equipment must be operated according to the specific conditions stated in the official three star rating for that equipment as published on HSE Chemicals Regulation Directorate's website.

Maintain three star operating conditions until 30 m from the top of the bank of any surface water bodies.

Aim spray away from water.

NOTE: BUFFER ZONES OF MORE THAN 5 M CANNOT BE REDUCED UNDER THE LOCAL ENVIRONMENT RISK ASSESSMENT FOR PESTICIDES (LERAP) SCHEME.

The statutory buffer zone must be maintained and the distance recorded in Section A of the LERAP record form.

The LERAP record form must be kept available for three years.

Storage and disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS. KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

EMPTY CONTAINER COMPLETELY and return to supplier.

PROTECT FROM FROST

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times.

Add washings to sprayer at time of filling and dispose of safely.

This label is compliant with the CPA Voluntary Initiative Guidance

5 L

Supplied by:

BASF plc, 4th and 5th Floors, 2 Stockport Exchange, Railway Road, Stockport SK1 3GG, Telephone: 0161 475 3000

**Emergency Information (24 hours freephone): 0049 180 227 3112
Technical Enquiries: 0845 602 2553 (office hours)**

® = Registered trademark of BASF

81165557 GB 2112



**The
Voluntary
Initiative**
16/1/2023

Pistol®

Warning

Very toxic to aquatic life with long lasting effects.

If medical advice is needed, have product container or label at hand.

Keep out of reach of children.

Read label before use.

Collect spillage.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

Contains 1,2-Benzisothiazol-3(2H)-one. May produce an allergic reaction.

To avoid risks to human health and the environment, comply with the instructions for use.



IMPORTANT INFORMATION

FOR USE ONLY AS AN INDUSTRIAL HERBICIDE

Situation: For use on natural surfaces not intended to bear vegetation, permeable surfaces overlaying soil, and hard surfaces (railway ballast only).

Maximum individual dose: 4.5 litres product/hectare

Maximum number of treatments: 1 per year

Latest time of application: End of July

Other specific restrictions:

This product must only be used on natural or porous surfaces such as gravel where the user can establish that the underlying surface is soil and railway ballast surfaces. Do not use if an impermeable membrane lies between the porous surface and the soil.

Do not apply to any non-porous man-made surfaces

Hand-held equipment: To minimise spray drift, the product must be applied using a nozzle capable of producing a coarse quality spray.

Vehicle-mounted equipment: Respect an unsprayed buffer zone to surface water bodies of 6 m. Respect an unsprayed buffer zone to surface water bodies of 6 m.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

GENERAL INFORMATION

PISTOL® is a suspension concentrate formulation containing 250 g/L glyphosate and 40 g/L diflufenican. PISTOL® has foliar absorbed translocated activity, combined with soil acting properties, providing pre-emergence weed control. The product is applied post-emergence to control a wide range of annual and perennial broad-leaved weeds and grasses on porous surfaces on land not intended to bear vegetation.

BASF Tip: For established deep rooted perennial weeds such as dandelion, thistle, dock and nettles which may re-generate, a post-emergence application is ideally required.

Restrictions/Warnings

Since there is a risk to aquatic life from use, direct spray from hand-held equipment must not be allowed to fall within 2 m of the top of the bank of any static or flowing waterbody or the top of a ditch which is dry at the time of application. Do not allow direct spray from horizontal boom sprayer to fall within 6 m of the top of the bank of a static or flowing water body, or within 1 m of the top of a ditch which is dry at the time of application. Spray must be aimed away from water.

Applications should not be made to plants growing under stress conditions, such as drought or water-logging, as reduced levels of control may result.

Do not spray in windy weather.

Extreme care must be taken to avoid drift onto non-target plants. This includes leaf surfaces, young bark or suckers of valued plants. Failure to do so may result in permanent damage or plant death.

Where PISTOL® has been applied to sites that are subsequently to be cleared or grubbed, a period of at least 6 months should elapse between treatment and the sowing and planting of subsequent crops. In addition, the soil should be ploughed or dug afterwards to ensure thorough mixing in order to remove any risk of damaging subsequent crops or planting.

Where PISTOL® or other products containing diflufenican are applied in successive years, levels of diflufenican will build up in the soil. Even where soils are thoroughly dug there may be a risk of damage to subsequent plantings.

Care should be taken when applying PISTOL® as heavy rain following application may wash the herbicide onto sensitive areas such as newly sown grass and areas about to be planted.

Where the soil organic matter content is greater than 10%, or for example where leaves have collected or where a mat of organic matter has built up, pre-emergence activity will be reduced.

For maximum persistence of activity the area treated should not be cultivated or raked following application.

For maximum pre-emergence and residual activity from PISTOL[®], ensure good coverage of the spray swath. Untreated areas will not be protected.

Overdosing should be avoided.

Do not add any wetting agent or adjuvant oil as this may result in reduced activity. However if using a CDA applicator, then an appropriate carrier may be used.

DO NOT APPLY PISTOL[®] OVER DRAINS OR IN DRAINAGE CHANNELS, GULLIES OR SIMILAR STRUCTURES FOR MOVING WATER.

WEEDS CONTROLLED

Strains of some annual weeds (e.g. black-grass, wild-oats, and Italian rye-grass) have developed resistance to herbicides which may lead to poor control. A strategy for preventing and managing such resistance should be adopted. This should include integrating herbicides with a programme of cultural control measures. Guidelines have been produced by the Weed Resistance Action Group and copies are available from the HGCA, CPA, your distributor, crop adviser or product manufacturer.

PISTOL[®] has a broad spectrum of activity if used at the recommended rate and under optimum conditions. It should give initial 'knockdown' of a wide variety of broad-leaved and grass weeds and thereafter residual control for a wide variety of broad-leaved weeds for at least 3 months, and up to 4 months on hard surfaces.

Weeds controlled such as:

Annual meadow-grass, annual nettle, buttercup, Canadian fleabane, mouse ear chickweed, chickweed, common daisy, common groundsel, ragwort, couch grass, cranesbill sp, dandelion, dock, fat hen, field forget-me-not, mayweed, perennial thistle, plantain sp, polygonaceae sp, red dead-nettle, rough-stalked meadow-grass, sowthistle, speedwell, willowherb. Effectiveness using three star drift reduction technology may be reduced.

Grasses may be the first weeds to return in treated areas and may require re-treatment with a glyphosate based product such as CDA Vanquish[®] Biactive.

BASF Tip: Treated areas which may have been protected from the spray swath such as areas of soil shaded by a larger weed canopy at application may require re-treatment with a glyphosate based product such as CDA Vanquish[®] Biactive.

Long term control

Perennial weeds such as dandelions and docks emerging from established root-stocks after application will not be controlled.

SITUATION SPECIFIC INFORMATION

Use Areas

PISTOL® may be used in non-crop areas against weeds in open soil and against weeds growing in gravel or other porous surfaces. Examples of suitable use areas include fence lines, strips of land adjacent to buildings, industrial sites, timber yards, farm yards, oil and gas storage sites, power stations, electric sub-stations, beneath pylons, around street/park obstacles.

PISTOL® may be used on railway track, railway sidings and other ballast areas of rail infrastructure.

PISTOL® must not be used on non-porous man made surfaces, for example paved areas, concrete or tarmac car parks and footpaths.

PISTOL® may be used on porous surfaces such as gravel ONLY where the underlying surface is soil. Do not use if an impermeable membrane lies between the gravel and the soil. The product must not be used on gravel where the underlying surface is concrete, tarmac or any other non-porous surface, or in situations where there is potential for run-off into surface waters.

Crop Specific Information

Rate of Use

EQUIPMENT	AREA	PRODUCT REQUIRED	WATER VOLUME	Spray Quality (Nozzle)	Spray height	Buffer Zone
Knapsack with 'conventional' nozzles	100m ²	45 ml	2 - 5 L	Coarse *Hypro Polijet AN0.6 or similar (provided)	40 cm	2 m
	1000 m ²	450 ml	20-50 L			
	1 ha	4.5 L	200-500 L			
Knapsack with CDA Applicator**	100m ²	45 ml	200 ml	Volume Median Diameter at least 300 microns	-	2 m
	1000 m ²	450 ml	2.0 L			
	1 ha	4.5 L	20.5 L			
Vehicle-mounted boom	1 ha	4.5 L	200-500 L	Three star drift reduction nozzles	-	6 m

*Hypro Polijet AN0.6 nozzle provided gives, subject to calibration: a coarse spray with a flow rate of 0.6 L/min at 1 bar, giving 225 L/ha at 4kph walking speed, swath width of 40 cm, 40 cm nozzle height

** Carrier volume and spray quality determined using a RiverLynx Eliminator CDA applicator. Use a carrier volume such that the total volume is equivalent to 25 L/ha.

Application Timing

Apply PISTOL® post-emergence of the weeds at any time from February until the end of July, provided that the weeds are actively growing and have not begun the process of natural die-back. At least six hours, but preferably 24 hours of dry weather are required immediately following application for optimum control.

BASF Tip: For post-emergence application ideally annual broad-leaved weeds should have at least two fully expanded true leaves and annual grasses should be at the one leaf growth stage or beyond. Some perennial weeds, including docks, perennial sowthistle and willowherb are best treated just before flowering or the setting of seed.

Mixing and Application

Mixing

Shake well before use. Half fill the spray tank with clean water (or carrier for CDA applicators). Start gentle agitation and add the required quantity of PISTOL®. Mix and top up to the required volume with water or carrier. Use immediately.

WASH OUT THE SPRAYER THOROUGHLY AFTER USE, USING A WETTING AGENT OR PROPRIETARY TANK CLEANER WITH TWO RINSES, AS TRACES OF PISTOL® MAY CAUSE HARM TO OTHER SUSCEPTIBLE PLANTS SPRAYED LATER.

Equipment

Hand-held equipment

Use a knapsack sprayer fitted with a coarse nozzle using a pressure of around 1-2 bars to provide a coarse spray. Use of anti-drift nozzles or the use of a protector shield to avoid any drift is recommended.

NOZZLE: Hypro Polijet AN0.6 nozzle (provided) gives, subject to calibration: a coarse spray with a flow rate of 0.6 L/min at 1 bar, giving 225L/ha at 4kph walking speed, swath width of 40 cm, 40 cm nozzle height, or use similar nozzles that give coarse spray.

Alternatively, Pistol may be used through specialist CDA equipment with a suitable droplet spectrum (VMD minimum 300 microns). Refer to the lance manufacturer's instructions before use.

Vehicle-mounted equipment

Apply via a horizontal boom sprayer. Apply in 200 – 500 L/ha using three star drift reducing nozzles. Use the higher volume where weed foliage is dense. A spray pressure of at least 2 bars is advised. Good, even spray coverage of soil and weeds is essential. To prevent damage, extreme care must be taken to avoid drift onto non-target plants. Good and even coverage of foliage and soil is essential for optimum activity.

Trade Mark Acknowledgments

PISTOL® is a registered trademark of BASF.

CDA VANQUISH® BIACTIVE is a registered trademark of Bayer CropScience Limited

The following does not form part of the authorised label text.

With many products there is a general risk of resistance developing to the active ingredients. For this reason a change in activity cannot be ruled out. It is generally impossible to predict with certainty how resistance may develop because there are so many crop and use connected ways of influencing this. We therefore have to exclude liability for damage or loss attributable to any such resistance that may develop. To help minimise any loss in activity the BASF recommended rate should in all events be adhered to.

Numerous, particularly regional or regionally attributable, factors can influence the activity of the product. Examples include weather and soil conditions, crop plant varieties, crop rotation, treatment times, application amounts, admixture with other products, appearance of organisms resistant to active ingredients and spraying techniques. Under particular conditions a change in activity or damage to plants cannot be ruled out. The manufacturer or supplier is therefore unable to accept any liability in such circumstances. All goods supplied by us are of high grade and we believe them to be suitable, but as we cannot exercise control over their mixing or use or the weather conditions during and after application, which may affect the performance of the material, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use; but nothing should be deemed to exclude or restrict any liability upon us which cannot be excluded or restricted under the provisions of the Unfair Contract Terms Act 1977 or any similar applicable law.

Section 6 of the Health and Safety at Work Act

Additional Product Safety Information

The product label provides information on a specific pesticidal use of the product; do not use otherwise, unless you have assessed any potential hazard involved, the safety measures required and that the particular use has “off-label” approval or is otherwise permitted under the Plant Protection Product Regulations (EC) No 1107/2009.

The information on this label is based on the best available information including data from test results.

Safety Data Sheet

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



agricentre.basf.co.uk/Pisto/MSDS

Alternatively, contact your supplier.

Pistol®



MAPP 18854

A suspension concentrate containing 250g/L glyphosate and 40g/L diflufenican.

A non-selective herbicide for control of annual and perennial weeds on natural surfaces not intended to bear vegetation, permeable surfaces overlaying soil and hard surfaces (railway ballast only).

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces, or applying the product.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

WASH HANDS AND EXPOSED SKIN before meals and after work.

Environmental protection

Do not contaminate water with the product.

Do not clean application equipment near surface water.

Avoid contamination via drains from farmyards and roads.

Extreme care must be taken to avoid spray drift onto non-crop plants outside the target area.

Hand-held equipment

Since there is a risk to aquatic life from use, direct spray must not be allowed to fall within 2 m of the top of the bank of any static or flowing waterbody or the top of a ditch which is dry at the time of application. Spray must be aimed away from water.

Vehicle-mounted equipment (boom sprayer)

To protect aquatic organisms, respect an unsprayed buffer zone to surface water bodies of 6 m.

HORIZONTAL BOOM SPRAYERS MUST BE FITTED WITH THREE STAR DRIFT REDUCTION TECHNOLOGY.

Low drift spraying equipment must be operated according to the specific conditions stated in the official three star rating for that equipment as published on HSE Chemicals Regulation Directorate's website.

Maintain three star operating conditions until 30 m from the top of the bank of any surface water bodies.

Aim spray away from water.

NOTE: BUFFER ZONES OF MORE THAN 5 M CANNOT BE REDUCED UNDER THE LOCAL ENVIRONMENT RISK ASSESSMENT FOR PESTICIDES (LERAP) SCHEME.

The statutory buffer zone must be maintained and the distance recorded in Section A of the LERAP record form.

The LERAP record form must be kept available for three years.

Storage and disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS. KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place. EMPTY CONTAINER COMPLETELY and return to supplier.

PROTECT FROM FROST

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times.

Add washings to sprayer at time of filling and dispose of safely.

This label is compliant with the CPA Voluntary Initiative Guidance

5 L

Supplied by:

BASF plc, 4th and 5th Floors, 2 Stockport Exchange, Railway Road, Stockport SK1 3GG, Telephone: 0161 475 3000

Emergency Information (24 hours freephone): 0049 180 227 3112

Technical Enquiries: 0845 602 2553 (office hours)

® = Registered trademark of BASF

81165557 GB 2112

