

■ BASF

We create chemistry

Pyrafen®

**MAPP 19913**

A suspension concentrate containing 250 g/l pyraclostrobin and 150 g/l boscalid. Pyrafen is a protectant and systemic fungicide for use against diseases in winter and spring oilseed rape.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

SAFETY PRECAUTIONS**Operator protection**

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

IF SWALLOWED, seek medical advice immediately and show this container or label.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH HANDS before meals and after work.

Environmental protection

Do not contaminate water with the product or its container.

Do not clean application equipment near surface water.

Avoid contamination via drains from farmyards and roads.

To protect aquatic organisms respect an unsprayed buffer zone to surface water bodies as specified for the crop.

HORIZONTAL BOOM SPRAYERS MUST BE FITTED WITH THREE STAR DRIFT REDUCTION TECHNOLOGY.

Low drift spraying equipment must be operated according to the specific conditions stated in the official three star rating for that equipment as published on HSE Chemicals Regulation Directorate's website.

Maintain three star operating conditions until 30 m from the top of the bank of any surface water bodies.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within the distance specified for the crop to the top of the bank of a static or flowing waterbody, or within 1m of the top of a ditch which is dry at the time of application.

Aim spray away from water.

NOTE: BUFFER ZONES OF MORE THAN 5 M CANNOT BE REDUCED UNDER THE LOCAL ENVIRONMENT RISK ASSESSMENT FOR PESTICIDES (LERAP) SCHEME.

The statutory buffer zone must be maintained and the distance recorded in Section A of the LERAP record form.

The LERAP record form must be kept available for three years.

Storage and disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

STORE IN ORIGINAL CONTAINER tightly closed, in a safe place.

DO NOT RE-USE CONTAINER for any purpose.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times.

Add washings to sprayer at time of filling and dispose of safely. Store in a suitable pesticide store. Keep dry and protect from frost.

This label is compliant with the CPA Voluntary Initiative Guidance

5 L

Supplied by: BASF plc, 4th and 5th Floors, 2 Stockport Exchange, Railway Road, Stockport SK1 3GG, Telephone: 0161 475 3000

**Emergency Information (24 hours freephone): 0049 180 227 3112
Technical Enquiries: 0845 602 2553 (office hours)**

® = Registered trademark of BASF

81171510 GB 2103

**The
Voluntary
Initiative**

25/10/2023

Pyrafen®

A suspension concentrate containing 250 g/l pyraclostrobin and 150 g/l boscalid.

Warning

Harmful if swallowed or inhaled.

Causes skin irritation.

May cause an allergic skin reaction.

May cause respiratory irritation.

Very toxic to aquatic life with long lasting effects.

Wear protective gloves/clothing/eye protection.

If medical advice is needed, have product container or label at hand.

Use only outdoors or in a well-ventilated area.


Call a POISON CENTRE or doctor/physician if you feel unwell.

Store in a well-ventilated place. Keep container tightly closed.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

Contains: 1,2-benzisothiazol-3(2H)-one (CAS 2634-33-5) and methylisothiazolinone (CAS 2682-20-4), may produce an allergic reaction.

To avoid risks to human health and the environment, comply with the instructions for use.



IMPORTANT INFORMATION

FOR USE ONLY AS A PROFESSIONAL FUNGICIDE, as directed below:

<u>Crop</u>	<u>Maximum individual dose</u>	<u>Maximum total dose per crop</u>	<u>Latest time of application</u>
Oilseed rape (winter and spring)	0.8 litre product/ha	0.8 litre product/ha	50% of pods have reached final size (GS 75)

Aquatic buffer zone distance: 12 metres

Other specific restrictions:

When used on crops with a greater than 5 m aquatic buffer zone, this product must not be applied via hand-held equipment.

Applications to oilseed rape (winter) must only be made between GS 14-75 – four leaves unfolded / 50% of pods have reached final size. Applications must not be made between 1 June and 31 October.

Applications to oilseed rape (spring) must only be made between GS 25-75 – five side shoots detectable / 50% of pods have reached final size. Applications must not be made between 1 June and 31 October.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

UFI: 64S3-J07K-6007-3G6Y

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Pyrafen is a protectant and systemic fungicide for use in winter and spring oilseed rape for the control of *Phoma lingam*/stem canker, Light leaf spot, Sclerotinia stem rot, and Alternaria dark leaf and pod spot.

Restrictions/Warnings

Avoid spray drift on to neighbouring crops and plants outside the target area.

Pyrafen represents minimal hazard to bees when used as directed. Consideration should be given to informing bee-keepers when you intend spraying crops in flower.

A reduction in Sclerotinia control may be observed when high-risk infection conditions occur after flowering, leading to secondary spread of the disease through plant-to-plant contact, which may be more severe in lodged crops.

Disease control

Pyrafen controls Phoma leaf spot and Stem canker, Sclerotinia stem rot, Alternaria dark leaf and pod spot and controls moderate infections of Light leaf spot in oilseed rape.

Disease	Susceptibility ratings
<i>Phoma</i> leaf spot #	Control
Stem canker #	Control
Light leaf spot #	Moderate control
Sclerotinia stem rot	Control
Alternaria dark leaf and pod spot	Moderate Control

Optimum control of Phoma leaf spot/stem canker and Light leaf spot will be achieved where Pyrafen is applied in the autumn/late winter or in the spring in sequence with an approved fungicide having activity on these diseases.

Resistance Management

GROUP

7

11

FUNGICIDES

Pyrafen contains boscalid, which belongs to the SDHI (carboxamide) group of fungicides and pyraclostrobin, which belongs to the strobilurin QoI group of fungicides. The two active ingredients possess different modes of action, which form part of a resistance management strategy.

The key to resistance management strategies is the reduction of selection pressure. This can be achieved by good agricultural practice which leads to less infection pressure (e.g. crop and stubble hygiene, use of less susceptible varieties, crop rotation, use of healthy seeds etc.).

Risk assessments for the target diseases on oilseed rape indicate a low or medium risk of resistance development. To minimize any possible risk of resistance occurring, Pyrafen should be applied in a preventative manner, following the recommendations on the label, and as an integral part of a spray program which includes fungicides with different modes of action.

For further advice on resistance management in oilseed rape contact your agronomist or specialist advisor and visit the Fungicide Resistance Action Committee (FRAC) and UK Fungicide Resistance Action Group (FRAG) websites.

Crop Specific Information

Pyrafen can be used on all varieties of winter and spring oilseed rape.

Time of application

Inspect crops to assess disease development prior to spraying. Apply Pyrafen protectively or when disease thresholds have been reached according to appropriate disease support systems.

Phoma leaf spot, Stem canker

Apply Pyrafen in the autumn at the first sign of disease. Optimum control will be achieved where Pyrafen is applied in sequence with an approved fungicide having activity on these diseases.

Light leaf spot

Apply Pyrafen protectively in the late autumn or at first sign of disease in early spring. Optimum control will be achieved where Pyrafen is applied in sequence with an approved fungicide having activity on these diseases.

Sclerotinia stem rot

Apply Pyrafen in high disease risk situations at early to full flower to coincide with or just before first petal fall, and before disease symptoms are visible. Applications made after the end of flowering (GS 69 BBCH) may result in reduced levels of Sclerotinia control.

Alternaria dark leaf and pod spot

Apply Pyrafen from full flowering up to and including when 50% of pods have reached final size. Applications made after the end of flowering may result in reduced levels of control.

Rate of application

Apply Pyrafen at 0.8 l/ha in a recommended water volume of 150-300 litres of water per hectare.

Mixing and Application

Mixing

Mixing procedure

Never prepare more spray solution than is required.

Three quarters fill the tank with clean water and start the agitation. To ensure thorough mixing of the product, invert the container several times before opening. Add the required quantity of Pyrafen to the spray tank while re-circulating. Add the remainder of the water and continue agitation until spraying is complete.

When tank mixes are to be used, take due note of any instructions given as to the order of mixing. Each product should be added separately to the spray tank and fully dispersed before the addition of any further product(s).

Rinse empty containers thoroughly, using an integrated pressure rinsing device or by manually rinsing three times. Add washings to tank at time of filling and dispose of container safely.

Application

Apply 0.8 l/ha Pyrafen in a recommended water volume of 150-300 litres of water per hectare as a MEDIUM spray, as defined by the BCPC. In dense crops, using the increased water volume will help achieve adequate penetration and good coverage of the crop.

Effectiveness using three star drift reduction technology may be reduced.

Where possible use narrow-wheeled high clearance application equipment. To ensure optimum spray coverage and minimize spray drift, adjust the spray boom to the appropriate height above the crop, according to guidance provided by the sprayer and/or nozzle manufacture.

Tank cleaning

After spraying, thoroughly clean and flush out application machinery with a minimum of three rinses.

Following Crops

Any crop can follow normally harvested or failed crops treated with Pyrafen.

Compatibility

For details of compatibilities contact your distributor, local BASF representative, the BASF Technical Services Hotline: 0845 602 2553 or visit our website on: www.agriCentre.basf.co.uk

Trade Mark Acknowledgments

Pyrafen[®] is a registered trademark of BASF.

Company Advisory Information

The following does not form part of the authorised label text.

With many products there is a general risk of resistance developing to the active ingredients. For this reason a change in activity cannot be ruled out. It is generally impossible to predict with certainty how resistance may develop because there are so many crop and use connected ways of influencing this. We therefore have to exclude liability for damage or loss attributable to any such resistance that may develop. To help minimise any loss in activity the BASF recommended rate should in all events be adhered to.

Numerous, particularly regional or regionally attributable, factors can influence the activity of the product. Examples include weather and soil conditions, crop plant varieties, crop rotation, treatment times, application amounts, admixture with other products, appearance of organisms resistant to active ingredients and spraying techniques. Under particular conditions a change in activity or damage to plants cannot be ruled out. The manufacturer or supplier is therefore unable to accept any liability in such circumstances. All goods supplied by us are of high grade and we believe them to be suitable, but as we cannot exercise control over their mixing or use or the weather conditions during and after application, which may affect the performance of the material, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us

for any damage or injury whatsoever arising from their storage, handling, application or use; but nothing should be deemed to exclude or restrict any liability upon us which cannot be excluded or restricted under the provisions of the Unfair Contract Terms Act 1977 or any similar applicable law.

Section 6 of the Health and Safety at Work Act

Additional Product Safety Information

The product label provides information on a specific pesticidal use of the product; do not use otherwise, unless you have assessed any potential hazard involved, the safety measures required and that the particular use has “off-label” approval or is otherwise permitted under the Plant Protection Product Regulations (EC) No 1107/2009.

The information on this label is based on the best available information including data from test results.

Safety Data Sheet

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



agricentre.basf.co.uk/Pyrafen/MSDS

Alternatively, contact your supplier.

EASY CONNECT



The easyconnect system requires a coupler for its use.

Step by step instruction for use with coupler:

- **Do NOT unscrew the easyconnect cap**
- Remove the dust cover
- **Do NOT push the plug manually into the container!**
- Turn the container upside down to connect the easyconnect cap to the coupler, then follow the instructions from your coupler supplier.
- After complete emptying and rinsing, push the plug into the container using a spike on the draining rack, allowing any residual rinsing water to drain.
- The easyconnect cap can be recycled together with the container according to local regulations. Do not transfer cap onto any other containers.
- **Note:** Whilst using the easyconnect coupler, please continue to wear PPE in line with the guidance on the product label.



Please find more information on www.easyconnect.tech

If you don't have a coupler yet, just unscrew the cap for conventional filling.

Step by step instruction for conventional use:

- **Do NOT remove the dust cover or push the plug into the container!**
- In the absence of a coupler, easyconnect caps can be used in the same way as traditional caps.
- Unscrew the easyconnect cap and pour the liquid into the induction bowl.
- **Note:** This will not leverage the benefits of the easyconnect system.
- The easyconnect cap can be recycled together with the container according to local regulations.

BASF

We create chemistry

Pyrafen[®]

**MAPP 19913**

A suspension concentrate containing 250 g/l pyraclostrobin and 150 g/l boscalid. Pyrafen is a protectant and systemic fungicide for use against diseases in winter and spring oilseed rape.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

SAFETY PRECAUTIONS**Operator protection**

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

IF SWALLOWED, seek medical advice immediately and show this container or label.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH HANDS before meals and after work.

Environmental protection

Do not contaminate water with the product or its container.

Do not clean application equipment near surface water.

Avoid contamination via drains from farmyards and roads.

To protect aquatic organisms respect an unsprayed buffer zone to surface water bodies as specified for the crop.

HORIZONTAL BOOM SPRAYERS MUST BE FITTED WITH THREE STAR DRIFT REDUCTION TECHNOLOGY.

Low drift spraying equipment must be operated according to the specific conditions stated in the official three star rating for that equipment as published on HSE Chemicals Regulation Directorate's website.

Maintain three star operating conditions until 30 m from the top of the bank of any surface water bodies.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within the distance specified for the crop to the top of the bank of a static or flowing waterbody, or within 1m of the top of a ditch which is dry at the time of application.

Aim spray away from water.

NOTE: BUFFER ZONES OF MORE THAN 5 M CANNOT BE REDUCED UNDER THE LOCAL ENVIRONMENT RISK ASSESSMENT FOR PESTICIDES (LERAP) SCHEME.

The statutory buffer zone must be maintained and the distance recorded in Section A of the LERAP record form.

The LERAP record form must be kept available for three years.

Storage and disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

STORE IN ORIGINAL CONTAINER tightly closed, in a safe place.

DO NOT RE-USE CONTAINER for any purpose.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times.

Add washings to sprayer at time of filling and dispose of safely. Store in a suitable pesticide store. Keep dry and protect from frost.

This label is compliant with the CPA Voluntary Initiative Guidance

® = Registered trademark of BASF

81171510 GB 2103

**The
Voluntary
Initiative****5 L**

Supplied by: BASF plc, 4th and 5th Floors, 2 Stockport Exchange, Railway Road, Stockport SK1 3GG, Telephone: 0161 475 3000

Emergency Information (24 hours freephone): 0049 180 227 3112
Technical Enquiries: 0845 602 2553 (office hours)