

BASF

We create chemistry

Tevos®

**MAPP 19789****An emulsifiable concentrate containing 142.5 g/litre pyraclostrobin and 75 g/litre fluxapyroxad.****A systemic fungicide for use in winter barley, spring barley, triticale, rye and oats.****The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.****SAFETY PRECAUTIONS****Operator protection**

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate and contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

Do not apply via hand-held equipment.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH HANDS AND EXPOSED SKIN before meals and after work.

Environmental protection

Do not contaminate water with the product or its container.

Do not clean application equipment near surface water.

Avoid contamination via drains from farmyards and roads. To protect aquatic organisms respect an unsprayed buffer zone to surface water bodies in line with LERAP requirements.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 metres of the top of the bank of a static or flowing water body, unless a Local Environment

Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 metre from the top of a ditch which is dry at the time of application. Aim spray away from water.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for inspection for three years.

Storage and disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times.

Add washings to sprayer at time of filling and dispose of safely.

Store in a suitable pesticide store. Keep dry and protect from frost.

DO NOT RE-USE CONTAINER FOR ANY PURPOSE.

This label is compliant with the CPA Voluntary Initiative Guidance

Supplied by:

BASF plc, 4th and 5th Floors, 2 Stockport Exchange, Railway Road, Stockport SK1 3GG, Telephone: 0161 475 3000

Emergency Information (24 hours freephone):

0049 180 227 3112

Technical Enquiries: 0845 602 2553 (office hours)



® = Registered trademark of BASF

81171822 GB 2103



**The
Voluntary
Initiative**
10/10/2023

5 L

Tevos®

An emulsifiable concentrate containing 142.5 g/litre pyraclostrobin and 75 g/litre fluxapyroxad.

Warning

Harmful if swallowed.

Harmful if inhaled.

May cause respiratory irritation.

May cause harm to breast-fed children.

Very toxic to aquatic life with long lasting effects.

Use only outdoors or in a well-ventilated area.

If medical advice is needed, have product container or label at hand.

IF INHALED: Remove person to fresh air and keep comfortable for breathing.

IF exposed or concerned: Call a POISON CENTRE or doctor/physician.


Store in a well-ventilated place. Keep container tightly closed.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

Contains propanoic acid, 2-hydroxy-, 2-ethylhexyl ester, (2S)-.

May produce an allergic reaction

To avoid risks to human health and the environment, comply with the instructions for use.



IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE, as directed below:

Crops	Maximum individual dose	Maximum number of applications	Latest time of application	Aquatic buffer zone distance (metres):
Winter and spring Barley, Rye, Triticale, Oats	1.5 litres product/ha	2 per crop	Up to and including Flowering (anthesis) just complete (BBCH 69)	5

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

UFI: 6KSM-S484-T00J-96HN

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Tevos is a protectant and systemic fungicide for use in winter barley, spring barley, triticale, rye and oats.

When applied for disease control this product may increase green leaf area retention of barley, rye and triticale.

Restrictions/Warnings

Extreme care must be taken to avoid drift onto nearby crops.

When applied to barley intended for malting, refer to the rate and latest timing restrictions for fluxapyroxad and pyraclostrobin given by the British Beer and Pub Association.

No more than two applications of any QoI fungicide may be made to crops of barley, oats, rye or triticale.

You must not apply more than two foliar applications of products containing complex II inhibitors (SDHIs) to any cereal crop.

After spraying, wash out sprayer thoroughly according to manufacturer's guideline and dispose of washings and clean containers according to DEFRA Code of Practice and local authority guidelines.

Disease control

Tevos is a protectant and systemic fungicide for the control of a range of diseases in winter barley, spring barley, triticale, rye and oats as summarised below:

	Winter barley	Spring barley	Oats	Rye	Triticale
<i>Septoria tritici</i>	-	-	-	-	C
Yellow rust	C	C	-	C	C
Brown rust	C	C	-	C	C
Crown rust	-	-	C	-	-
<i>Rhynchosporium</i>	C	C	-	C	-
Net blotch	C	C	-	-	-
<i>Ramularia</i>	C	C	-	-	-
Powdery mildew	MC	MC	MC	MC	MC

C = Control

MC = Moderate Control

When applied for disease control this product may increase green leaf area retention of barley, rye and triticale.

Resistance Management

GROUP	7	11	FUNGICIDES
-------	---	----	------------

Tevos should be used in accordance with the instructions for use for the target diseases at the specific growth stages indicated.

Use Tevos as part of an Integrated Crop Management (ICM) strategy incorporating other methods of control, including where appropriate other fungicides with a different mode of action.

Tevos contains pyraclostrobin, a member of the QoI cross resistance group.

Tevos should be used preventatively and should not be relied upon for its curative potential.

You must not apply more than two foliar applications of QoI containing products to any cereal crop.

Strains of barley powdery mildew resistant to QoIs are common in the UK.

Tevos contains fluxapyroxad which belongs to the SDHI (carboxamide) complex II inhibitor group of fungicides.

To minimise the risk of resistance occurring, apply a maximum of two SDHI fungicide sprays to any cereal crop.

There is a significant risk of widespread resistance occurring in *Septoria tritici* populations in the UK. Failure to follow resistance management action may result in reduced levels of disease control.

For the control of *Septoria* leaf blotch, Tevos must always be used in mixture with another product recommended for its control, and that contains a fungicide from a different cross resistance group to SDHIs and QoIs and is applied at a dose that will give robust control

Visit the FRAG-UK website for further advice on resistance management of cereal fungicides.

Crop Specific Information

Tevos is a protectant and systemic fungicide for use in winter barley, spring barley, triticale, rye and oats.

Time of Application

Apply Tevos at the start of foliar disease attack. A maximum of two applications can be made up to and including flowering just complete in winter and spring barley, oats and rye and triticale.

Earliest timing for first application: BBCH 25

Earliest timing for second application: BBCH 30

Following Crops

After treating a cereal crop with Tevos, cabbage, carrot, clover, dwarf French beans, field beans, lettuce, maize, oats, oilseed rape, onions, peas, potatoes, ryegrass, sugar beet, sunflowers, barley and wheat may be sown as the following crop. The effect of Tevos on other following crops has not been assessed.

Mixing and Application

Mixing

Half fill the spray tank with clean water and start the agitation. SHAKE THE CONTAINER WELL before use and pour in the required amount of product. Rinse any empty containers thoroughly and add rinsings to the spray tank. Add the remainder of the water and continue agitation until spraying is completed.

When tank mixes are to be used, each product should be added separately to the spray tank, taking due note of any instructions given as to the order of mixing.

Application

Apply 1.5 litres of Tevos per hectare in 150 to 300 litres of water per hectare.

Qualified recommendation

Tevos may be applied at 1.5 litres per hectare in 100 litres water per hectare, although efficacy at this reduced volume has not been evaluated. Therefore application at 100 litres water per hectare is at user's risk with regard to biological efficacy.

Apply as a **MEDIUM** spray, as defined by BCPC.

Compatibility

For details of compatibilities contact your distributor, local BASF representative, the BASF Technical Services Hotline: 0845 602 2553 or visit our website www.agriCentre.basf.co.uk

Trade Mark Acknowledgments

® = Registered trademark of BASF.

The following does not form part of the authorised label text.

With many products there is a general risk of resistance developing to the active ingredients. For this reason a change in activity cannot be ruled out. It is generally impossible to predict with certainty how resistance may develop because there are so many crop and use connected ways of influencing this. We therefore have to exclude liability for damage or loss attributable to any such resistance that may develop. To help minimise any loss in activity the BASF recommended rate should in all events be adhered to.

Numerous, particularly regional or regionally attributable, factors can influence the activity of the product. Examples include weather and soil conditions, crop plant varieties, crop rotation, treatment times, application amounts, admixture with other products, appearance of organisms resistant to active ingredients and spraying techniques. Under particular conditions a change in activity or damage to plants cannot be ruled out. The manufacturer or supplier is therefore unable to accept any liability in such circumstances. All goods supplied by us are of high grade and we believe them to be suitable, but as we cannot exercise control over their mixing or use or the weather conditions during and after application, which may affect the performance of the material, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use; but nothing should be deemed to exclude or restrict any liability upon us which cannot be excluded or restricted under the provisions of the Unfair Contract Terms Act 1977 or any similar applicable law.

Section 6 of the Health and Safety at Work Act

Additional Product Safety Information

The information on this label is based on the best available information including data from test results.

Safety Data Sheet

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



agricentre.basf.co.uk/Tevos/MSDS

Alternatively, contact your supplier.

BASF

We create chemistry

Tevos®

**MAPP 19789**

An emulsifiable concentrate containing 142.5 g/litre pyraclostrobin and 75 g/litre fluxapyroxad.

A systemic fungicide for use in winter barley, spring barley, triticale, rye and oats.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

SAFETY PRECAUTIONS**Operator protection**

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate and contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

Do not apply via hand-held equipment.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH HANDS AND EXPOSED SKIN before meals and after work.

Environmental protection

Do not contaminate water with the product or its container.

Do not clean application equipment near surface water.

Avoid contamination via drains from farmyards and roads.

To protect aquatic organisms respect an unsprayed buffer zone to surface water bodies in line with LERAP requirements.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 metres of the top of the bank of a static or flowing water body, unless a Local Environment

Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 metre from the top of a ditch which is dry at the time of application.

Aim spray away from water.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for inspection for three years.

Storage and disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times.

Add washings to sprayer at time of filling and dispose of safely.

Store in a suitable pesticide store. Keep dry and protect from frost.

DO NOT RE-USE CONTAINER FOR ANY PURPOSE.

This label is compliant with the CPA Voluntary Initiative Guidance

Supplied by:

BASF plc, 4th and 5th Floors, 2 Stockport Exchange, Railway Road, Stockport SK1 3GG, Telephone: 0161 475 3000

Emergency Information (24 hours freephone): 0049 180 227 3112

Technical Enquiries: 0845 602 2553 (office hours)



® = Registered trademark of BASF

81171822 GB 2103



The
Voluntary
Initiative

5 L