

Dymid®



Product Registration Number: MAPP 17585

A suspension concentrate containing 500 g/litre propyzamide and 5.3 g /litre aminopyralid (present as 6.3 g/litre aminopyralid potassium salt).

A foliar and residual herbicide for the control of a wide range of weeds in WINTER OILSEED RAPE.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

SAFETY PRECAUTIONS

Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate or handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WASH HANDS before meals and after work.

Environmental protection:

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads. Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target area.

Storage and disposal:

STORE IN ORIGINAL CONTAINER, tightly closed, in a safe place.

EMPTY CONTAINER COMPLETELY and dispose of safely.

Protect from frost.

10 L

This label is compliant with the
CPA Voluntary Initiative Guidance



Approval Holder:
Dow AgroSciences Limited
CPC2 Capital Park, Fulbourn,
Cambridge, CB21 5XE.
Telephone: (01462) 457272
Fax: (01462) 426605
24 Hour Emergency Telephone Number:
+44 (0) 1553 761 251

Marketing Company:
BASF plc
Crop Protection
PO BOX 4 Earl Road,
Cheadle Hulme, CHEADLE,
Cheshire, SK8 6QG
Tel.: +44 (0)161 485 6222
24 Hour Emergency Telephone Number:
+49 180 2273112
Technical Enquiries:
0845 602 2553 (office hours)


We create chemistry

® = Trademark of the Dow Chemical Company ("Dow") or an affiliated company of Dow

P99053811810

81130577GB1118



Dymid®

Product Identifier according to Art.18 of Reg. (EC) No 1272/2008 [CLP]:
Dymid®, Propyzamide

Warning:
Suspected of causing cancer.
Very toxic to aquatic life with long lasting effects.

Do not handle until all safety precautions have been read and understood.
Wear protective gloves/protective clothing/eye protection/face protection.
IF exposed or concerned: Get medical advice/attention.
Collect spillage.
Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean, triple rinsed containers which can be disposed of as non-hazardous waste.

Contains 1,2-benzisothiazolin-3(2H)-one. May produce an allergic reaction.

To avoid risks to human health and the environment, comply with the instructions for use.



IMPORTANT INFORMATION.

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

Crops/Situations	Maximum Individual Dose (litres product/hectare)	Maximum Number of Treatments	Latest Time of Application
Oilseed rape (winter)	1.7	One per crop	Before 1st February in year of harvest

Other Specific Restrictions:

Do not harvest crops for human or animal consumption for at least 6 weeks after application.
Livestock must be kept out of treated areas for at least 1 week following treatment and until poisonous weeds such as ragwort have died and become unpalatable.
Users must have received adequate instruction, training and guidance in the safe and efficient use of the product and must take all reasonable precautions to protect the health of human beings and non-target organisms and safeguard the environment.
The product must not be used on land where vegetation will be cut for animal feed, fodder or bedding nor for composting or mulching within one calendar year of treatment.
Following crops (winter and spring wheat) should not be planted within 30 weeks of application of **DYMID**.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

IMPORTANT INFORMATION.

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

Crops/Situations	Maximum Individual Dose (litres product/hectare)	Maximum Number of Treatments	Latest Time of Application
Oilseed rape (winter)	1.7	One per crop	Before 1st February in year of harvest

Other Specific Restrictions:

Do not harvest crops for human or animal consumption for at least 6 weeks after application.

Livestock must be kept out of treated areas for at least 1 week following treatment and until poisonous weeds such as ragwort have died and become unpalatable.

Users must have received adequate instruction, training and guidance in the safe and efficient use of the product and must take all reasonable precautions to protect the health of human beings and non-target organisms and safeguard the environment.

The product must not be used on land where vegetation will be cut for animal feed, fodder or bedding nor for composting or mulching within one calendar year of treatment.

Following crops (winter and spring wheat) should not be planted within 30 weeks of application of **DYMID**.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

WARNINGS

DYMID contains aminopyralid; traces of this material in oilseed rape straw can damage sensitive crops. Aminopyralid residues in plant tissues which have not completely decayed may affect succeeding susceptible crops e.g. peas, beans and other legumes, sugar beet, fodder beet, carrots and umbelliferae, potatoes and tomatoes, lettuce and other compositae. Therefore following good agricultural practice ensure that all remains of the oil seed rape crop have completely decayed before planting susceptible crops.

DO NOT remove oilseed rape straw from the field unless it is to be used for the burning for heat or electricity production.

DO NOT feed animals with treated oilseed rape straw.

DO NOT Use oilseed rape straw for animal bedding.

DO NOT Use oilseed rape straw for composting or mulching.

Take care to avoid local overdosing.

Do not make more than one application of DYMID to any one crop.

If animals are inadvertently fed or bedded on oilseed rape straw treated with DYMID, do not use animal waste for composting.

Do not use on crops grown for seed.

Neighbouring Crops/Plants

Avoid damage by drift onto susceptible crops, non-target plants or waterways. Do not apply directly to, or allow spray drift to come into contact with agricultural or horticultural crops, amenity plantings, gardens, ponds, lakes or watercourses.

SOIL TYPES

DYMID can be used on all soil types except those containing more than 10% organic matter.

SOIL AND WEATHER CONDITIONS

DYMID requires moisture for root uptake. Best residual action is obtained in moist soils of fine tilth.

DYMID can be applied under frosty conditions but should not be used where run off from the soil surface is likely.

Best results are achieved when growth of weeds (especially blackgrass and volunteer cereals) is slow, but transpiration continues.

In mild autumns/winters, emerged weeds may take longer to be controlled, the residual activity of DYMID will be shortened and overall control may be reduced.

The efficacy of DYMID may be reduced in organic soils and in the presence of excessive surface organic debris, burnt straw, ash, or ploughed up turf.

RESISTANCE

Strains of some annual grasses (e.g. blackgrass, wild oats, Italian ryegrass) have developed resistance to herbicides which may lead to poor control. A strategy for preventing and managing such resistance should be adopted. Guidelines have been produced by the Weed Resistance Action Group and copies are available from the HGCA, CPA, your distributor, crop adviser or product manufacturer.

WINTER OILSEED RAPE

DYMID can be applied after the use of an approved specific graminicide applied in accordance with the manufacturer's recommendations.

WATER VOLUMES

Apply DYMID in 200 to 300 litres of water per hectare.

Ensure good ground cover.

APPLICATION EQUIPMENT

Application should be made through a ground crop sprayer.

Do not apply through broadcast air-assisted sprayers.

CROP RECOMMENDATION TABLES

S = Susceptible MS = Moderately susceptible MR = Moderately resistant R = Resistant

Crop	Rate of Use	Weed Species	Stage of Weed Growth			Time of Year	Timing Stage of Crop	Soil Type (Soil Texture (85 System))
			Germinating	Up to 2 leaf	Established*			
Winter oilseed rape	1.5 L/ha	Annual meadow grass, barren brome, volunteer cereals, wild oat	S	S	S	1st October to 31st January	As soon as possible after 3rd true leaf stage. Crop selectivity is by depth protection. Factors which cause shallow rooting may reduce crop selectivity.	All soils with less than 10% organic matter.
		Common chickweed', mayweed, Common poppy	S	S	S'			
		Blackgrass ²	S	S	MS ²			
		Black bindweed, black nightshade, fat hen, knotgrass, redshank, small nettle, speedwells	S	S	MR			
		Field forget me not	MS	MS	R			
		Cleavers	MS	MR	R			
	1.0 L/ha - Use this rate only where a specific graminicide has controlled volunteer cereals and grass weeds and chickweed is not a problem	Annual meadow grass, volunteer cereals, wild oat	S	S	S			
		Black bindweed, black nightshade, common chickweed, fat hen, knotgrass, redshank, small nettle	S	S	R			
		Speedwells	S	R	R			

* * Established = 3-4 true leaves to flower bud development stage.

NOTES FOR CROP RECOMMENDATION TABLES

1	Chickweed control may be reduced where it is well established (over 10 cm in diameter)
2	<p>CONTROL OF BLACKGRASS Established (well tillered) blackgrass is moderately susceptible: Where populations of blackgrass and/or volunteer cereals exceed 50/m² DYMID should be applied in tank mix with an approved graminicide, or following an effective approved graminicide to ensure optimum weed control. Deeper germinating blackgrass within the soil profile could reduce product efficacy. Where partial resistance (R# or RR#) to the partner graminicide is known to exist the dose of DYMID may be increased to 1.7 L/ha. This may also be done if applications are made early in the season, under warm conditions and an increase in the duration of residual control is required (see 'Soil and Weather Conditions' above). Where high levels of resistance (RRR#) to the partner graminicide occurs there is no advantage of adding this graminicide to DYMID for blackgrass control, and DYMID even at 1.7 L/ha will not give acceptable levels of established black-grass control in these circumstances. # R = 1* RR = 2*/3* RRR = 4*/5*</p>

FOLLOWING CROPS

Winter cereals and spring cereals only.

Treated land must be mouldboard ploughed to a depth of 15 cm prior to drilling a following cereal crop.

Please consult Dow AgroSciences if a treated crop fails because of bad growing conditions.

MIXING

Add half the required volume of water to the spray tank and begin agitation. Add the recommended quantity of DYMID. Agitate while topping up the tank and continue agitation until spraying is complete.

TANK CLEANING

Thoroughly wash all spraying and measuring equipment with water immediately after use.

To avoid subsequent injury to crops, all spraying equipment must be thoroughly cleaned both inside and out after an application of DYMID.

1. Immediately after spraying, drain tank completely. Any contamination on the outside of the spraying equipment should be removed by washing with clean water.
2. Rinse inside of tank with clean water and flush through booms and hoses using at least one tenth of the spray tank volume. Drain tank completely.
3. Half fill tank with clean water. Agitate and then briefly flush the boom and hoses. Top up with water making sure the tank is completely full and allow to stand for 15 minutes with agitation. Flush the boom and hoses and drain tank completely.
4. Nozzles and filters should be cleaned separately and removed if necessary.
5. For disposal of washings, follow local regulations. Do not spray onto sensitive crop or land intended for cropping with sensitive crop.

Note: If it is not possible to drain the tank completely, step 3 must be repeated before going onto step 4.

TRADEMARK ACKNOWLEDGEMENTS

DYMID is a trademark of the Dow Chemical Company (Dow) or an affiliate company of Dow.

All other brand names used in this document are trademarks of other manufacturers in which proprietary rights may exist.

Dow AgroSciences Conditions of Supply

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use, or the weather conditions before, during or after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded. No responsibility will be accepted by us or re sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

**Section 6 of the Health and Safety at Work Act
Additional Product Safety Information**

The product label provides information on a specific pesticidal use of the product; do not use otherwise, unless you have assessed any potential hazard involved, the safety measures required and that the particular use has "off-label" approval or is otherwise permitted under the Plant Protection Product Regulations (EC) No 1107/2009.

The information on this label is based on the best available information including data from test results

Safety Data Sheet

To access the Safety Data Sheet for this product scan the QR code or use the weblink below.



http://www.agricentre.basf.co.uk/go/dymid_sds

Alternatively, contact your supplier.

Dymid®



Product Registration Number: MAPP 17585

A suspension concentrate containing 500 g/litre propyzamide and 5.3 g /litre aminopyralid (present as 6.3 g/litre aminopyralid potassium salt).

A foliar and residual herbicide for the control of a wide range of weeds in WINTER OILSEED RAPE.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

SAFETY PRECAUTIONS

Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate or handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WASH HANDS before meals and after work.

Environmental protection:

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads. Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target area.

Storage and disposal:

STORE IN ORIGINAL CONTAINER, tightly closed, in a safe place.

EMPTY CONTAINER COMPLETELY and dispose of safely.

Protect from frost.

10 L

This label is compliant with the
CPA Voluntary Initiative Guidance



Approval Holder:

Dow AgroSciences Limited
CPC2 Capital Park, Fulbourn,
Cambridge, CB21 5XE.

Telephone: (01462) 457272

Fax: (01462) 426605

24 Hour Emergency Telephone Number:
+44 (0) 1553 761 251

Marketing Company:

BASF plc
Crop Protection
PO BOX 4 Earl Road,
Cheadle Hulme, CHEADLE,
Cheshire, SK8 6QG

Tel.: +44 (0)161 485 6222

24 Hour Emergency Telephone Number:
+49 180 2273112

Technical Enquiries:

0845 602 2553 (office hours)

 **BASF**
We create chemistry

® = Trademark of the Dow Chemical Company ("Dow") or an affiliated company of Dow

P99053811810

81130577GB1118