

Zampro® DM

MAPP No: 15013

A suspension concentrate containing

300 g/l ametoctradin and 225 g/l dimethomorph.

A fungicide for use in potatoes.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

SAFETY PRECAUTIONS

Operator protection

WHEN USING DO NOT EAT, DRINK OR SMOKE.

IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

Environmental protection

Do not contaminate surface waters or ditches with chemical or used container.

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

Storage and disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

DO NOT RE-USE CONTAINER for any purpose.

STORE IN ORIGINAL CONTAINER tightly closed, in a safe place.

On emptying the container, RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of container safely.

5 L

This label is compliant with the
CPA Voluntary Initiative Guidance



Supplied by:

BASF plc

Agricultural Solutions

PO Box 4, Earl Road

Cheadle Hulme, CHEADLE

Cheshire SK8 6QG

Tel: 0161 485 6222

Emergency information,

freephone 24 hrs:

0049 180 2273 112

Technical Enquiries:

0845 602 2553 (office hours)

® = Registered trademark of BASF

 **- BASF**
We create chemistry

81130304GB1118



Zampro® DM

A suspension concentrate containing
300 g/l ametoctradin and 225 g/l dimethomorph

**WARNING:
HARMFUL IF SWALLOWED
VERY TOXIC TO AQUATIC LIFE
WITH LONG LASTING EFFECTS**

WASH WITH PLENTY OF WATER AND SOAP
THOROUGHLY AFTER HANDLING.
DO NOT EAT, DRINK OR SMOKE WHEN USING THIS
PRODUCT.

IF SWALLOWED: CALL A POISON CENTRE OR DOCTOR/
PHYSICIAN IF YOU FEEL UNWELL.
RINSE MOUTH.

DISPOSE OF CONTENTS/CONTAINER TO A
LICENSED HAZARDOUS-WASTE DISPOSAL
CONTRACTOR OR COLLECTION SITE EXCEPT
FOR EMPTY CLEAN CONTAINERS WHICH CAN BE
DISPOSED OF AS NON-HAZARDOUS WASTE.

**To avoid risks to human health and the environment,
comply with the instructions for use.**



This product is approved under the Plant Protection Products Regulations.

IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE, as directed below:

Crop	Maximum Individual dose	Maximum total dose	Latest time of application
Potato	0.8 litres product per hectare	3.2 litres product per hectare	7 days before harvest

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Zampro DM is a fungicide with a combination of protectant, translaminar and locally systemic leaf activity for the control of foliar late blight (*Phytophthora infestans*) and the reduction of tuber blight in potatoes.

1. Disease Control

Zampro DM will control foliar late blight on leaves and reduce the incidence of tuber blight when used as part of a full programme of correctly timed sprays.

1.1 Disease susceptibility

Zampro DM is most effective when applied as a protectant spray before the risk of blight occurs.

Zampro DM will give effective protection against phenylamide resistant strains of late blight.

1.2 Resistance management

Zampro DM contains a mixture of ametoctradin (a triazolo-pyrimidylamine, Qxl fungicide) and dimethomorph (a CAA fungicide).

To minimize the risk of resistance developing, and in line with current FRAG-UK advice to use effective doses, use Zampro DM at the recommended rate (0.8 l/ha). No more than 3 consecutive applications of Zampro DM, and no more than a total of 4 applications of Zampro DM per season should be applied.

General advice:

Up to 6 applications of a CAA fungicide in mixture with a fungicide with a different mode of action, making up no more than half of the total number of intended late blight sprays per season, may be made.

No more than three consecutive applications of a CAA fungicide should be made.

Up to 6 applications of a Qxl fungicide in mixture with a fungicide with a different mode of action, making up no more than half of the total number of intended late blight sprays per season, may be made.

No more than three consecutive applications of a Qxl fungicide should be made.

Zampro DM will control strains of late blight resistant to phenylamides. No cross resistance to phenylamide resistant strains of blight have been observed.

If an intensive blight control programme is necessary, include products with different modes of action. When fungicides with the same mode of action are used repeatedly over several years in the same field, naturally occurring less sensitive strains may survive, propagate and become dominant in that field. It is therefore advised that wherever possible spray programmes should include fungicides with differing modes of action. Zampro DM contains active substances with two different modes of action. For further advice on resistance management for potato blight (*Phytophthora infestans*) contact your agronomist or specialist advisor and/or visit the FRAG-UK and Potato Council websites.

2. Crop specific information

2.1 Crops

Zampro DM can be used on all varieties of potato, including seed crops.

2.2 Time of application

Foliar blight

Zampro DM should be applied as part of a repeat spray programme at 7–10 day intervals, depending on disease pressure. As disease pressure and the risk of late blight infection increases, the interval should be shortened. Irrigated crops should be treated as high risk.

Applications of Zampro DM should begin as soon as there is a risk of blight infection, or an official blight warning has been issued. In the absence of any warnings, applications should commence before the crop meets along the rows.

Tuber blight

To reduce the incidence of tuber blight at harvest, Zampro DM should be used at the recommended dose and spray intervals from tuber initiation to complete haulm desiccation.

2.3 Rate of application

0.8 l/ha.

3. Mixing and Spraying

3.1 Mixing

Half fill the sprayer tank with clean water and start the agitation. Add the required amount of Zampro DM. Add the remainder of the water and continue agitation until spraying is completed. Never prepare more spray solution than is required.

3.2 Application

Apply as a medium spray, as defined by BCPC, in a water volume of 200–500 l/ha. Select the appropriate water volume for the density of the crop, ensuring good coverage of the foliage and stems.

Zampro DM is rainfast in 1 hour, providing application has been made to dry foliage and the spray has dried on the leaf.

3.3 Tank cleaning

Thoroughly wash out spray equipment after use.

The following does not form part of the product label under the Plant Protection Products Regulations.

With many products there is a general risk of resistance developing to the active ingredients. For this reason a change in activity cannot be ruled out. It is generally impossible to predict with certainty how resistance may develop because there are so many crop and use connected ways of influencing this. We therefore have to exclude liability for damage or loss attributable to any such resistance that may develop. To help minimise any loss in activity the BASF recommended rate should in all events be adhered to.

Numerous, particularly regional or regionally attributable, factors can influence the activity of the product. Examples include weather and soil conditions, crop plant varieties, crop rotation, treatment times, application amounts, admixture with other products, appearance of organisms resistant to active ingredients and spraying techniques. Under particular conditions a change in activity or damage to plants cannot be ruled out. The manufacturer or supplier is therefore unable to accept any liability in such circumstances. All goods supplied by us are of high grade and we believe them to be suitable, but as we cannot exercise control over their mixing or use or the weather conditions during and after application, which may affect the performance of the material, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use; but nothing should be deemed to exclude or restrict any liability upon us which cannot be excluded or restricted under the provisions of the Unfair Contract Terms Act 1977 or any similar applicable law.

**Section 6 of the Health and Safety at Work Act
Additional Product Safety Information**

The product label provides information on a specific pesticidal use of the product; do not use otherwise, unless you have assessed any potential hazard involved, the safety measures required and that the particular use has "off-label" approval or is otherwise permitted under the Plant Protection Products Regulation.

The information on this label is based on the best available information including data from test results.

Safety Data Sheet

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



bit.ly/Zampro_DM_sds

Alternatively, contact your supplier.