



Caryx®



MAPP 16100

A water-based formulation containing 30 g/l metconazole and 210 g/l mepiquat-chloride.

A plant growth regulator for use in winter oilseed rape.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

IN CASE OF CONTACT WITH EYES, RINSE IMMEDIATELY with plenty of water and seek medical advice.

Environmental protection

Do not contaminate water with the product or its container.

Do not clean application equipment near surface water.

Avoid contamination via drains from farmyards and roads.

To protect aquatic organisms respect an unsprayed buffer zone distance to surface water bodies in line with LERAP requirements.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 metres of the top of the bank of a static or flowing water body, unless a Local Environmental Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 metre from the top of a ditch which is dry at the time of application.

Aim spray away from water.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained.

The results of the LERAP must be recorded and kept available for three years.

Storage and disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

STORE IN ORIGINAL CONTAINER tightly closed, in a safe place. **RINSE CONTAINER THOROUGHLY** by using an integrated pressure rinsing device or manually rinsing three times.

Add washings to sprayer at time of filling and dispose of container safely.

This label is compliant with the CPA Voluntary Initiative Guidance



© Registered trademark of BASF

811 36023 GB 1079



5 L

Supplied by: BASF plc, Agricultural Solutions
PO Box 4, Earl Road, Cheadle Hulme, CHEADLE,
Cheshire SK8 6QG, Tel: 0161 485 6222

Emergency Information: (24 hours freephone): 0049 180 2273112
Technical Enquiries: 0845 602 2553 (office hours)



Caryx®

A water-based formulation containing 30 g/l metconazole and 210 g/l mepiquat-chloride.

Danger


Harmful if swallowed.
May cause an allergic skin reaction.
Causes serious eye damage.
Harmful if inhaled.
Very toxic to aquatic life with long lasting effects.

Wear protective gloves and eye/face protection.
Do not breathe mist or vapour.
IF ON SKIN (or hair): Wash with plenty of soap and water.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

Immediately call a POISON CENTER or doctor/physician.
Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.



IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL PLANT GROWTH REGULATOR, as directed below:

Crop	Timing	Maximum Individual Dose (litres product/ha)	Maximum Number of Treatments (per crop)	Latest Time of Application
Winter oilseed rape	Spring (GS 31-59) Beginning of stem extension to yellow bud	1.4	1	Up to and including crop growth stage 59 (Yellow bud)
	Sequence Autumn GS 13-19 followed by Spring GS 31-59	1.4	1	Up to and including crop growth stage 59 (Yellow bud)
		1.4	1	

Other specific restrictions:

This product must not be applied via hand-held equipment.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

CARYX is a plant growth regulator for the reduction of crop height and lodging risk in all varieties of winter oilseed rape.

By reducing the incidence and/or severity of lodging CARYX may help to protect from yield losses associated with lodged oilseed rape crops

Restrictions/Warnings

Avoid boom overlaps and ensure that the boom height is correctly adjusted.

Extreme care should be taken to avoid damage by drift onto plants outside the target area.

Ensure the sprayer is free from previous chemical residues that may harm the crop. Use of a detergent cleaner is advised before and after use.

Use of a growth regulation product can offer good protection from the incidence or severity of lodging but there can be no guarantee that lodging will not occur, especially in high lodging pressure situations.

Crop Specific Information

CARYX can be used on all varieties of winter oilseed rape.

Application of CARYX

Spring application of CARYX

CARYX applied at 1.4 l/ha in the spring will shorten the crop and manipulate canopy structure, reducing the risk of the crop lodging.

Application timing

Apply CARYX in the spring between crop growth stage 31-59 of the crop (beginning of stem extension to yellow bud) once the crop is actively growing. Applications made before the crop has shown signs of active growth will be less effective.

Reduction of lodging risk

CARYX applied in the spring will reduce the risk of lodging later in the season.

Maintaining the rate of CARYX will provide a more consistent reduction of lodging risk by lowering the incidence or severity of lodging.

Autumn followed by spring sequence of CARYX

Caryx may be used in an autumn / spring sequence to regulate crop growth going into winter; and shorten the crop and manipulate canopy structure in the spring, reducing the risk of lodging later in the season.

Use of this sequence is only recommended on forward crops.

Application timing

Apply Caryx as 1.4 l/ha in the autumn between growth stages 14-19 of the crop (4-9 leaves) and follow up with a second application of 1.4 l/ha in the spring between growth stages 31-59 of the crop (beginning of stem extension to yellow bud).

Autumn applications should only be made to forward crops. A forward crop can be defined as a crop that has the potential to grow to more than 25 cm tall by the **end** of autumn growing season. Shorter crops will benefit less from an autumn application and are best managed by solo spring applications.

The potential for a crop to reach this height is dependent on a number of factors for example; weather conditions post application, fertiliser regime, plant population, the nutrient and water content of the soil and variety.

Both autumn and spring applications should only be made when crops are actively growing and conditions remain conducive to continued growth.

Applications made after crop growth has started to slow in the autumn or before the crop has shown signs of active growth in the spring will be less effective.

Following Crops

Following crops after normally harvested winter oilseed rape

After treating a crop of oilseed rape with CARYX, the crops listed below may be sown as a following crop:-

- Beans
- Clover
- Oilseed rape
- Ryegrass
- Cabbage
- Lettuce
- Onions
- Sugar beet
- Carrots
- Linseed
- Peas
- Sunflower
- Cereals
- Maize
- Potatoes

The effects of Caryx on other crops has not been assessed.

Re-drilling due to crop failure in the autumn

In the event of crop failure, the following crops may be sown after cultivating to a depth of at least 15 cm:

Re-drilling the same autumn:

Winter wheat (excluding durum wheat), winter barley, winter oats, winter field beans, winter oilseed rape.

Re-drilling the following spring:

Spring cereals (including oats), maize, peas, beans, spring oilseed rape, linseed, sugar beet, sunflower, onions.

Crops not suitable for re-drilling after crop failure in the autumn

Due to a risk of crop damage it is recommended that grass crops, for example ryegrass, are not re-drilled in the same autumn or subsequent spring following a crop failure in the autumn.

Mixing and Application

Mixing

Never prepare more spray solution than is required.

SHAKE WELL BEFORE USE.

Fill the spray tank three quarters full with water and start the agitation.

To ensure thorough mixing of the product, invert the container several times before opening.

Add the required quantity of CARYX to the spray tank while re-circulating.

Fill the tank with the remaining required amount of water and continue agitation until spraying is completed.

On emptying the product container, rinse container thoroughly using an integrated pressure rinsing device or by manually rinsing three times.

Add washings to sprayer at time of filling and dispose of container safely.

Application

Apply CARYX in 200-400 l/ha water as a MEDIUM spray, as defined by BCPC.

Qualified Approval:

CARYX may be applied at up to 1.4 l/ha in 150 litres water per hectare although efficacy at this reduced volume has not been evaluated. Therefore application at 150 litres water per hectare is at user's risk with regard to biological efficacy.

To ensure optimum spray coverage and minimize spray drift, adjust the spray boom to the appropriate height above the crop according to guidance provided by the sprayer and/or nozzle manufacture.

Tank cleaning

Wash sprayer thoroughly immediately after use, using clean water and following the sprayer cleaning guidance provided by the equipment manufacturer.

Compatibility

For details of compatibilities contact your distributor, local BASF representative or the BASF Technical Services Hotline: 0044 845 602 2553

The following does not form part of the authorised label text.

With many products there is a general risk of resistance developing to the active ingredients. For this reason a change in activity cannot be ruled out. It is generally impossible to predict with certainty how resistance may develop because there are so many crop and use connected ways of influencing this. We therefore have to exclude liability for damage or loss attributable to any such resistance that may develop. To help minimise any loss in activity the BASF recommended rate should in all events be adhered to.

Numerous, particularly regional or regionally attributable, factors can influence the activity of the product. Examples include weather and soil conditions, crop plant varieties, crop rotation, treatment times, application amounts, admixture with other products, appearance of organisms resistant to active ingredients and spraying techniques. Under particular conditions a change in activity or damage to plants cannot be ruled out. The manufacturer or supplier is therefore unable to accept any liability in such circumstances. All goods supplied by us are of high grade and we believe them to be suitable, but as we cannot exercise control over their mixing or use or the weather conditions during and after application, which may affect the performance of the material, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use; but nothing should be deemed to exclude or restrict any liability upon us which cannot be excluded or restricted under the provisions of the Unfair Contract Terms Act 1977 or any similar applicable law.

Section 6 of the Health and Safety at Work Act

Additional Product Safety Information

The product label provides information on a specific pesticidal use of the product; do not use otherwise, unless you have assessed any potential hazard involved, the safety measures required and that the particular use has "off-label" approval or is otherwise permitted under the Plant Protection Products Regulations.

The information on this label is based on the best available information including data from test results.

Safety Data Sheet

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



bit.ly/Caryx_sds

Alternatively, contact your supplier.

BASF

We create chemistry

Caryx®



MAPP 16100

A water-based formulation containing 30 g/l metconazole and 210 g/l mepiquat-chloride.

A plant growth regulator for use in winter oilseed rape.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

IN CASE OF CONTACT WITH EYES, RINSE IMMEDIATELY with plenty of water and seek medical advice.

Environmental protection

Do not contaminate water with the product or its container.

Do not clean application equipment near surface water.

Avoid contamination via drains from farmyards and roads.

To protect aquatic organisms respect an unsprayed buffer zone distance to surface water bodies in line with LERAP requirements.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 metres of the top of the bank of a static or flowing water body, unless a Local Environmental Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 metre from the top of a ditch which is dry at the time of application.

Aim spray away from water.

This product qualifies for inclusion within the Local Environmental Risk Assessment for Pesticides (LERAP) scheme.

Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained.

The results of the LERAP must be recorded and kept available for three years.

Storage and disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

STORE IN ORIGINAL CONTAINER tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times.

Add washings to sprayer at time of filling and dispose of container safely.

This label is compliant with the CPA Voluntary Initiative Guidance



© Registered trademark of BASF

81136023 GB 1079



5 L

Supplied by: BASF plc, Agricultural Solutions
PO Box 4, Earl Road, Cheadle Hulme, CHEADLE,
Cheshire SK8 6QG, Tel: 0161 485 6222

Emergency Information: (24 hours freephone): 0049 180 2273112
Technical Enquiries: 0845 602 2553 (office hours)

