

BASF

We create chemistry

Enervin[®] SC

MAPP 19095

**A suspension concentrate containing 200 g/l ametoctradin.
A fungicide for use in potatoes and leeks.**

**The (COSHH) Control of Substances Hazardous to Health
Regulations may apply to the use of this product at work.**

SAFETY PRECAUTIONS

Operator protection

**WHEN USING DO NOT EAT, DRINK OR
SMOKE.**

**IF YOU FEEL UNWELL, seek medical
advice immediately (show the label
where possible).**

Environmental protection

**Do not contaminate water with the
product or its container.**

**Do not clean application equipment
near surface water.**

**Avoid contamination via drains from
farmyards and roads.**

Storage and disposal

**KEEP AWAY FROM FOOD, DRINK AND
ANIMAL FEEDING STUFFS.**

KEEP OUT OF REACH OF CHILDREN.

**DO NOT RE-USE CONTAINER for any
other purpose.**

**STORE IN ORIGINAL CONTAINER
tightly closed, in a safe place.**

**On emptying the container, RINSE
CONTAINER THOROUGHLY by using an
integrated pressure rinsing device or
manually rinsing three times.**

**Add washings to sprayer at time of
filling and dispose of container safely.**

**This label is compliant with the
CPA Voluntary Initiative Guidance**

5 L

Supplied by: BASF plc, Agricultural Solutions
PO Box 4, Earl Road, Cheadle Hulme, CHEADLE,
Cheshire SK8 6QG, Tel: 0161 485 6222

Emergency Information: (24 hours freephone): 0049 180 2273112,
Technical Enquiries: 0845 602 2553 (office hours)

® = Registered trademark of BASF

81131207 GB 1049



**The
Voluntary
Initiative**

Enervin® SC

A suspension concentrate containing 200 g/l ametoctradin.

Toxic to aquatic life with long lasting effects.

If medical advice is needed, have product container or label at hand.

Keep out of reach of children.

Collect spillage.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

May produce an allergic reaction. Contains 1,2-BENZISOTHAZOL-3(2H)-ONE.

To avoid risks to human health and the environment, comply with the instructions for use.



IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE, as directed below:

Crop	Maximum Individual dose	Maximum total dose	Latest time of application
Potato	1.2 litres product per hectare	4.8 litres product per hectare	7 days before harvest
Leek	1.2 litres product per hectare	2.4 litres product per hectare	7 days before harvest

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Enervin SC is a protectant fungicide for use in the early stages of foliar late blight (*Phytophthora infestans*) in potato and white tip (*Phytophthora porri*) in leek.

Disease control

Enervin SC will provide protection against the early stages of foliar late blight in potato and white tip in leek. A full programme of correctly timed sprays using other authorised fungicides is likely to be required for longer term control and particularly where disease pressure is high.

Disease susceptibility

Enervin SC is most effective when applied as a protectant spray before the risk of late blight in potato or white tip in leek occurs.

Enervin SC will give effective protection against phenylamide resistant strains of late blight.

Resistance Management

Enervin SC contains ametoctradin (a triazolo-pyrimidylamine, QoSI fungicide).

To minimize the risk of resistance developing, and in line with current FRAG-UK advice to use effective doses, always use Enervin SC at the recommended rate (1.2 l/ha) and in mixture with products with a different mode of action at a dose that will give robust control.

In potato no more than 3 consecutive applications of Enervin SC, and no more than a total of 4 applications of Enervin SC per season should be applied.

In leek no more than a total of 2 applications of Enervin SC per season should be applied.

No more than 6 applications of a QoSI fungicide, making up no more than half of the total number of intended late blight sprays per season, may be made.

Enervin SC should be used early in the spray programme, after which application with other fungicides should be made that can offer sufficient protection under high disease conditions. In this way sub-optimal control of *Phytophthora infestans* is prevented, thereby managing the resistance risk.

General advice:

In potato, Enervin SC will control strains of late blight resistant to phenylamides. No cross resistance to phenylamide resistant strains of blight have been observed.

If an intensive blight control programme is necessary, include products with different modes of action.

When fungicides with the same mode of action are used repeatedly over several years in the same field, naturally occurring less sensitive strains may survive, propagate and become dominant in that field. It is therefore advised that wherever possible spray programmes should include fungicides with differing modes of action. For further advice on resistance management for potato blight (*Phytophthora infestans*) contact your agronomist or specialist advisor and/or visit the FRAG-UK and Potato Council websites.

Crop Specific Information

Potatoes

Enervin SC can be used on all varieties of potato, including seed crops.

Time of application

Enervin SC should be applied as part of a repeat spray programme at 5 – 10 day intervals, depending on disease pressure. As disease pressure and the risk of late blight infection increases, the interval should be shortened. Irrigated crops should be treated as high risk. Applications of Enervin SC should begin as soon as there is a risk of blight infection, or an official blight warning has been issued. In the absence of any warnings, applications should commence before the crop meets along the rows.

Rate of application

1.2 L/ha Enervin SC in 200-400 L/ha of water.

Leek

Qualified recommendation for a minor use:

There is limited evidence of crop safety and effectiveness for the control of *Phytophthora porri* in leek

Time of application

Enervin SC should be applied as part of a repeat spray programme at 7 – 10 day intervals, depending on disease pressure. As disease pressure and the risk of white tip infection increases, the interval should be shortened. Irrigated crops should be treated as high risk.

Applications of Enervin SC should begin as soon as there is a risk of white tip.

Rate of application

1.2 L/ha Enervin SC in 350-1000 L/ha water.

Mixing and Application

Application

Apply in a water volume of 200 – 400 l/ha for potato and 350 – 1000 l/ha for leek, selecting the appropriate water volume for the density of the crop, ensuring good coverage of the foliage and stems.

Apply as a MEDIUM spray, as defined by BCPC.

Mixing

Never prepare more spray solution than is required.

Three quarters fill the tank with clean water and start the agitation. To ensure thorough mixing of the product, invert the container several times before opening. Add the required quantity of Enervin SC to the spray tank while re-circulating. Add the remainder of the water and continue agitation until spraying is complete.

When tank mixes are to be used, take due note of any instructions given as to the order of mixing. Each product should be added separately to the spray tank and fully dispersed before the addition of any further product(s).

All tank mixes should be used immediately after mixing.

Tank cleaning

Wash sprayer thoroughly immediately after use, using clean water and following the sprayer cleaning guidance provided by the equipment manufacturer.

Compatibility

For details of compatibilities contact your distributor, local BASF representative, the BASF Technical Services Hotline: 0845 602 2553 or visit our website on: www.agriCentre.basf.co.uk

The following does not form part of the authorised label text.

With many products there is a general risk of resistance developing to the active ingredients. For this reason a change in activity cannot be ruled out. It is generally impossible to predict with certainty how resistance may develop because there are so many crop and use connected ways of influencing this. We therefore have to exclude liability for damage or loss attributable to any such resistance that may develop. To help minimise any loss in activity the BASF recommended rate should in all events be adhered to.

Numerous, particularly regional or regionally attributable, factors can influence the activity of the product. Examples include weather and soil conditions, crop plant varieties, crop rotation, treatment times, application amounts, admixture with other products, appearance of organisms resistant to active ingredients and spraying techniques. Under particular conditions a change in activity or damage to plants cannot

be ruled out. The manufacturer or supplier is therefore unable to accept any liability in such circumstances. All goods supplied by us are of high grade and we believe them to be suitable, but as we cannot exercise control over their mixing or use or the weather conditions during and after application, which may affect the performance of the material, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use; but nothing should be deemed to exclude or restrict any liability upon us which cannot be excluded or restricted under the provisions of the Unfair Contract Terms Act 1977 or any similar applicable law.

Section 6 of the Health and Safety at Work Act

Additional Product Safety Information

The product label provides information on a specific pesticidal use of the product; do not use otherwise, unless you have assessed any potential hazard involved, the safety measures required and that the particular use has “off-label” approval or is otherwise permitted under the Plant Protection Products Regulations.

The information on this label is based on the best available information including data from test results.

Safety Data Sheet

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



bit.ly/Enervin_SC_sds

Alternatively, contact your supplier.

BASF

We create chemistry

Enervin[®] SC

MAPP 19095

**A suspension concentrate containing 200 g/l ametoctradin.
A fungicide for use in potatoes and leeks.**

**The (COSHH) Control of Substances Hazardous to Health
Regulations may apply to the use of this product at work.**

SAFETY PRECAUTIONS

Operator protection

WHEN USING DO NOT EAT, DRINK OR
SMOKE.

IF YOU FEEL UNWELL, seek medical
advice immediately (show the label
where possible).

Environmental protection

Do not contaminate water with the
product or its container.
Do not clean application equipment
near surface water.
Avoid contamination via drains from
farmyards and roads.

Storage and disposal

KEEP AWAY FROM FOOD, DRINK AND
ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

DO NOT RE-USE CONTAINER for any
other purpose.

STORE IN ORIGINAL CONTAINER
tightly closed, in a safe place.

On emptying the container, RINSE
CONTAINER THOROUGHLY by using an
integrated pressure rinsing device or
manually rinsing three times.

Add washings to sprayer at time of
filling and dispose of container safely.

**This label is compliant with the
CPA Voluntary Initiative Guidance**

5 L

Supplied by: BASF plc, Agricultural Solutions
PO Box 4, Earl Road, Cheadle Hulme, CHEADLE,
Cheshire SK8 6QG, Tel: 0161 485 6222

Emergency Information: (24 hours freephone): 0049 180 2273112,
Technical Enquiries: 0845 602 2553 (office hours)

® = Registered trademark of BASF

81131207 GB 1049



**The
Voluntary
Initiative**