

Environmental Information Sheet

PICOSTOMP MAFF/MAPP 13455



A suspension concentrate formulation containing 16 g/litre picolinafen (pyridinecarboxamide herbicide) and 320 g/litre pendimethalin (dinitroaniline herbicide) for use in winter wheat and winter barley.

Maximum application rate: 3 litres/hectare (48g + 960 g a.s./ha)

Maximum number of treatments: One per crop

Section	Profile
<p>1. WILDLIFE</p> <p>Mammals and Birds</p>	<p>PicoStomp is not classified as '<i>Harmful to game, wild birds and animals</i>'.</p> <p>No risk management is necessary to protect wild mammals and birds. Picolinafen is of low toxicity to birds and mammals. Pendimethalin is of low toxicity to mammals and moderate toxicity to birds. The risk to wild mammals and birds grazing on treated areas is low as is the risk due to exposure from other routes e.g. consumption of invertebrates such as insects.</p>
<p>2. BEES</p>	<p>No risk management necessary. PicoStomp is of low risk to bees.</p>
<p>3. NON TARGET INSECTS AND OTHER ARTHROPODS</p>	<p>No risk management necessary. PicoStomp poses a low risk to a range of arthropod species commonly found in and around treated fields, including ground beetles and spiders.</p>
<p>4. AQUATIC LIFE</p>	<p>PicoStomp is "<i>VERY TOXIC TO AQUATIC ORGANISMS, may cause long term adverse effects in the aquatic environment</i>".</p> <p>PicoStomp is of high toxicity to aquatic invertebrates e.g. water flea and fish and of extreme toxicity to green algae. '<i>Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads</i>'.</p> <p>Risk management is essential. PicoStomp can be used safely providing care is taken to prevent spray drift reaching surface waters. The following risk management practices must be carried out in order to ensure that there is adequate protection for aquatic species: '<i>DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing water body or within 1m of the top of a ditch which is dry at the time of application. Aim spray away from water</i>'.</p> <p>LERAP Category B. Buffer zones may be reduced (see LERAP Guidelines).</p>

Environmental Information Sheet

PICOSTOMP MAFF/MAPP 13455



Section	Profile
<p>5. SOIL and GROUNDWATER</p> <p>Earthworms</p> <p>Soil Micro-organisms</p>	<p>Picolinafen is moderately persistent and pendimethalin is very persistent in soil. Both picolinafen and pendimethalin are strongly bound to soil and can be considered not mobile. Use of PicoStomp according to the label presents a low risk to groundwater. No risk management necessary.</p> <p>Picolinafen and pendimethalin are of low toxicity to earthworms. No risk management necessary.</p> <p>No effects on soil microbial processes were found so the risk to soil micro-organisms is considered low. No risk management is necessary.</p>
<p>6. NON-TARGET PLANTS</p>	<p>PicoStomp is a herbicide for control of meadow grasses and broad-leaved weeds. <i>'Extreme care should be taken to avoid damage by drift on to plants outside the target area'</i>.</p> <p>Care should be taken to minimise drift of PicoStomp onto conservation headlands, beetle banks and field boundaries such as hedgerows, which may provide an important habitat for wild flowers.</p>

ALWAYS READ THE LABEL: USE PESTICIDES SAFELY

Care must be taken to minimise the risk of surface water contamination from farmyard and field sources.

For further information about the environmental profile of this product contact:-

BASF plc, Agricultural Division, PO Box 4, Earl Road, Cheadle Hulme, Cheadle, Cheshire SK8 6QG Telephone: 0161 485 6222 Fax: 0161 486 0891

This Environmental Information Sheet was prepared in accordance with CPA Guidance Notes Version 4.

© copyright of BASF plc.