

Document size: 115x130 mm (width x height)  
Gross size: 135x130 mm (width x height)



# Flyer® 200



MAPP 17293

**An emulsifiable concentrate containing 200 g/litre (19.2% w/w) pyraclostrobin. A protectant fungicide with curative properties, for use in winter wheat, winter barley, spring wheat, spring barley, oats.**

**A protectant fungicide for use in forage and grain maize.**

**The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.**

## SAFETY PRECAUTIONS

### Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

AVOID ALL CONTACT WITH SKIN.

WASH HANDS AND EXPOSED SKIN before meals and after work.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

### Environmental protection

Do not contaminate water with the product or its container.

Do not clean equipment near surface water.

Avoid contamination via drains from farmyards and roads.

To protect aquatic organisms respect an unsprayed buffer zone distance to surface water bodies in line with LERAP requirements.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 metres of the top of the bank of a static or flowing water body, unless a Local Environmental Risk

Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 metre from the top of a ditch which is dry at the time of application.

DO NOT ALLOW DIRECT SPRAY from hand-held sprayers to fall within 1 metre of the top of the bank of a static or flowing water body.

Aim spray away from water.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained.

The results of the LERAP must be recorded and kept available for inspection for three years.

### Storage and disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS. KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of container safely.

Do not re-use container for any purpose.

Store in a suitable pesticide store. Keep dry and protect from frost.

**This label is compliant with the CPA Voluntary Initiative Guidance**



© Registered trademark of BASF

81138391 GB 1119



# 5 L

Supplied by: BASF plc, Agricultural Solutions  
PO Box 4, Earl Road, Cheadle Hulme, CHEADLE,  
Cheshire SK8 6QG, Tel: 0161 485 6222

Emergency Information: (24 hours freephone): 0049 180 2273112  
Technical Enquiries: 0845 602 2553 (office hours)



## Flyer® 200

An emulsifiable concentrate containing  
200 g/litre (19.2% w/w) pyraclostrobin.

### Danger

**Harmful if swallowed.**  
**May be fatal if swallowed and enters airways.**  
**Causes skin irritation.**  
**May cause an allergic skin reaction.**  
**Causes serious eye irritation.**  
**Harmful if inhaled.**  
**Very toxic to aquatic life with long lasting effects.**

Wear protective gloves/clothing.  
Avoid breathing vapours.  
Wash with plenty of water and soap thoroughly after handling.

IF SWALLOWED: Immediately call a POISON CENTRE or doctor/physician.

IF ON SKIN: Wash with plenty of soap and water.

Do NOT induce vomiting.

**To avoid risks to human health and the environment, comply with the instructions for use.**



### IMPORTANT INFORMATION

**FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE, as directed below:**

Crops	Maximum individual dose	Maximum number of treatments	Latest time of application
Winter and spring wheat	1.25 litres product per hectare	2 per crop	Before grain watery ripe (GS 71)
Winter and spring barley, Oats	1.25 litres product per hectare	2 per crop	Up to and including emergence of ear just complete (GS 59)
Maize (Forage and Grain)	1.0 litre product per hectare	1 per crop	Up to and including mid flowering (BBCH 65)

Other specific restrictions:

The maximum concentration must not exceed 1.25 litres of product in 200 litres of water.

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**

## DIRECTIONS FOR USE

**IMPORTANT:** This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Flyer 200 is a fungicide with protectant and curative properties for disease control in winter and spring wheat, winter and spring barley and oats.

Flyer 200 is a fungicide with protectant properties for disease control in maize (forage and grain).

In cereals yield response may be obtained in the absence of visual disease.

In maize a grain yield response may be obtained in the absence of visual disease.

Flyer can maintain the total greenness of the crop towards the end of the growing season.

## Restrictions/Warnings

**Extreme care must be taken to avoid drift onto nearby crops.**

Wash equipment thoroughly after use.

You must not apply more than two foliar applications of QoI containing products to any cereal crop. See also Section, Resistance Management, below.

## Disease control

Flyer 200 is a fungicide with protectant and curative properties for disease control in winter and spring wheat, winter and spring barley and oats. Flyer 200 is a fungicide with protectant properties for disease control in maize (forage and grain). Details are summarised below:

	Winter Wheat	Spring Wheat	Winter Barley	Spring Barley	Oats	Maize (Forage and grain)
<i>Septoria tritici</i> (See Note 1)	C	C	-	-	-	-
<i>Septoria nodorum</i> (See Note 2)	MC	MC	-	-	-	-
Yellow rust	C	C	C	C	-	-
Brown rust	C	C	C	C	-	-
Crown rust	-	-	-	-	C	-
Net blotch	-	-	C	C	-	-
<i>Rhynchosporium</i>	-	-	MC	MC	-	-
Eyespot	-	-	-	-	-	C
Northern leaf blight	-	-	-	-	-	MC

C = Control

MC = Moderate Control

**NOTE 1:** *S. tritici*: Flyer 200 is recommended for control of *S. tritici* with moderate curative ability in the latent phase.

**NOTE 2:** *S. nodorum*: Flyer 200 is recommended for control of *S. nodorum* with best results achieved when used as a protective treatment or during early stages of disease establishment.

### Resistance Management

Flyer 200 contains pyraclostrobin, a member of the QoI cross resistance group. Flyer 200 should be used preventatively and should not be relied on for its curative potential.

On cereal crops, Flyer 200 must always be used in mixture with another product, recommended for control of the same target disease that contains a fungicide from a different cross resistance group and is applied at a dose that will give robust control.

Use Flyer 200 as part of an Integrated Crop Management (ICM) strategy incorporating other methods of control, including where appropriate other fungicides with a different mode of action. You must not apply more than two foliar applications of QoI containing products to any cereal crop.

There is a significant risk of widespread resistance occurring in *Septoria tritici* populations in the UK. Failure to follow resistance management action may result in reduced levels of disease control.

### Crop Specific Information

Flyer 200 is a fungicide with protectant and curative properties for disease control in winter and spring wheat, winter and spring barley and oats.

Flyer 200 is a fungicide with protectant properties for disease control in maize (forage and grain).

### Time of Application

#### Cereals

Apply Flyer 200 at the start of foliar disease attack. A maximum of two applications can be made before grain watery ripe in winter and spring wheat and up to and including emergence of ear just complete in winter and spring barley and oats.

#### Maize (Forage and Grain)

Best results are achieved by an application of Flyer 200 before the start of foliar disease attack or at the very earliest signs of infection.

Flyer 200 gives control of Eyespot for up to seven weeks after application when applied before the start of foliar disease attack.

If applying before the start of disease attack, it is recommended to only treat sites that have a known history of disease when conditions are conducive for disease development. It is recommended to consult with a specialist maize agronomist before application.

Once established in the crop Flyer 200 can reduce the development of the disease.

Application can be made up to and including mid-flowering.

## Rate of Application

### Cereals

Apply 1.25 litres Flyer 200 in a minimum of 200 litres of water per hectare.

### Maize (Forage and Grain)

Apply 1.0 litre Flyer 200 in 200-400 litres of water per hectare.

## Mixing and Application

### Mixing

Half fill the spray tank with clean water and start the agitation. SHAKE THE CONTAINER WELL before use and pour in the required amount of product. Rinse any empty containers thoroughly and add rinsings to the spray tank. Add the remainder of the water and continue agitation until spraying is completed. When tank mixes are to be used, each product should be added separately to the spray tank. All tank mixes should be used immediately after mixing.

### Application

Apply as a MEDIUM spray, as defined by BCPC.

### Compatibility

For details of compatibilities contact your distributor, local BASF representative or the BASF Technical Services Hotline: 0044 845 602 2553.

### The following does not form part of the authorised label text.

With many products there is a general risk of resistance developing to the active ingredients. For this reason a change in activity cannot be ruled out. It is generally impossible to predict with certainty how resistance may develop because there are so many crop and use connected ways of influencing this. We therefore have to exclude liability for damage or loss attributable to any such resistance that may develop. To help minimise any loss in activity the BASF recommended rate should in all events be adhered to. Numerous, particularly regional or regionally attributable, factors can influence the activity of the product. Examples include weather and soil conditions, crop plant varieties, crop rotation, treatment times, application amounts, admixture with other products, appearance of organisms resistant to active ingredients and spraying techniques. Under particular conditions a change in activity or damage to plants cannot be ruled out. The manufacturer or supplier is therefore unable to accept any liability in such circumstances. All goods supplied by us are of high grade and we believe them to be suitable, but as we cannot exercise control over their mixing or use or the weather conditions during and after application, which may affect the performance of the material, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use; but nothing should be deemed to exclude or restrict any liability upon us which cannot be excluded or restricted under the provisions of the Unfair Contract Terms Act 1977 or any similar applicable law.

## Section 6 of the Health and Safety at Work Act

### Additional Product Safety Information

The product label provides information on a specific pesticidal use of the product; do not use otherwise, unless you have assessed any potential hazard involved, the safety measures required and that the particular use has "off-label" approval or is otherwise permitted under the Plant Protection Product Regulations (EC) No 1107/2009.

The information on this label is based on the best available information including data from test results.

### Safety Data Sheet

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



[bit.ly/Flyer\\_200\\_sds](https://bit.ly/Flyer_200_sds)

Alternatively, contact your supplier.



# Flyer® 200



MAPP 17293

**An emulsifiable concentrate containing 200 g/litre (19.2% w/w) pyraclostrobin. A protectant fungicide with curative properties, for use in winter wheat, winter barley, spring wheat, spring barley, oats.**

**A protectant fungicide for use in forage and grain maize.**

**The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.**

## SAFETY PRECAUTIONS

### Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

**WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD)** when handling the concentrate.

**WEAR SUITABLE PROTECTIVE GLOVES** when handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

**AVOID ALL CONTACT WITH SKIN.**

**WASH HANDS AND EXPOSED SKIN** before meals and after work. **WHEN USING DO NOT EAT, DRINK OR SMOKE.**

**IF YOU FEEL UNWELL**, seek medical advice immediately (show the label where possible).

### Environmental protection

Do not contaminate water with the product or its container.

Do not clean equipment near surface water.

Avoid contamination via drains from farmyards and roads.

To protect aquatic organisms respect an unsprayed buffer zone distance to surface water bodies in line with LERAP requirements.

**DO NOT ALLOW DIRECT SPRAY** from horizontal boom sprayers to fall within 5 metres of the top of the bank of a static or flowing water body, unless a Local Environmental Risk Assessment for Pesticides (LERAP) permits a narrower buffer

zone, or within 1 metre from the top of a ditch which is dry at the time of application.

**DO NOT ALLOW DIRECT SPRAY** from hand-held sprayers to fall within 1 metre of the top of the bank of a static or flowing water body.

**Aim spray away from water.**

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme.

Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained.

The results of the LERAP must be recorded and kept available for inspection for three years.

### Storage and disposal

**KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS. KEEP OUT OF REACH OF CHILDREN.**

**KEEP IN ORIGINAL CONTAINER**, tightly closed, in a safe place.

**RINSE CONTAINER THOROUGHLY** by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of container safely.

Do not re-use container for any purpose.

Store in a suitable pesticide store. Keep dry and protect from frost.

**This label is compliant with the CPA Voluntary Initiative Guidance**



© Registered trademark of BASF

81138391 GB 1119



# 5 L

Supplied by: BASF plc, Agricultural Solutions  
PO Box 4, Earl Road, Cheadle Hulme, CHEADLE,  
Cheshire SK8 6QG, Tel: 0161 485 6222

Emergency Information: (24 hours freephone): 0049 180 2273112  
Technical Enquiries: 0845 602 2553 (office hours)