

# Belanty®

MAPP 20548

A suspension concentrate containing 75 g/l mefenentrifluconazole. An agricultural/horticultural fungicide for the control of a range of diseases in maize, sweetcorn, potatoes, combining pea, vining pea, field bean, beans without pods (dry), cabbage, Brussels sprout, broccoli/calabrese, cauliflower, carrot, parsnip, celeriac, swede and turnip.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

## SAFETY PRECAUTIONS

### Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

DO NOT HANDLE TREATED BRASSICA CROPS for 9 days post-treatment.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH HANDS AND EXPOSED SKIN before meals and after work.

IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

### Environmental protection

Do not contaminate water with the product or its container.

Do not clean application equipment near surface water.

Avoid contamination via drains from farmyards and roads.

### Storage and disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place, away from sources of heat.

DO NOT RE-USE CONTAINER FOR ANY PURPOSE.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times.

Add washings to sprayer at time of filling and dispose of container safely.

Store in a suitable pesticide store. Keep dry and protect from frost.

This label is compliant with the CPA Voluntary Initiative Guidance

#### Supplied by:

BASF plc, 4th and 5th Floors, 2 Stockport Exchange, Railway Road, Stockport, SK1 3GG  
Telephone: 0161 475 3000

Emergency Information (24 hours freephone):

0049 180 227 3112

Technical Enquiries: 0845 602 2553  
(office hours)

® = Registered trademark of BASF

81188901 GB 2026



The  
Voluntary  
Initiative  
5/2/2026

# Belanty®

A suspension concentrate containing 75 g/l  
mefentri fluconazole.

## Warning

**May cause an allergic skin reaction.  
Toxic to aquatic life with long lasting effects.**

Avoid breathing mist/vapours/spray.

Wear protective gloves/clothing.

Wash contaminated clothing before reuse.

Collect spillage.

If skin irritation or rash occurs: Call a POISON CENTRE or doctor/physician.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

Contains isocyanates [CAS No. 1887149-14-5] and isothialozones [BIT CAS No. 2634-33-5, CIT/MIT 3:1 CAS No. 55965-84-9 and MIT CAS No. 2682-20-4]. May produce an allergic reaction.

**To avoid risks to human health and the environment, comply with the instructions for use.**



## IMPORTANT INFORMATION

### FOR USE ONLY AS A PROFESSIONAL FUNGICIDE, as directed below:

Crops	Maximum individual dose	Maximum total dose	Latest time of application
Grain maize, Forage maize, Sweetcorn	1.25 l/ha	1.25 l/ha	End of flowering (BBCH 69)
Potatoes	1.25 l/ha	3.75 l/ha	3 days before harvest
Field beans, Beans without pods (dry), Combining pea, Vining pea	1.3 l/ha	2.6 l/ha	20% of pods have reached typical length (BBCH 72)
Carrot, Parsnip, Celeriac, Swede, Turnip	1.5 l/ha	4.5 l/ha	3 days before harvest
Cabbage, Cauliflower, Broccoli/calabrese, Brussels sprouts	1.5 l/ha	4.5 l/ha	14 days before harvest

### Other Specific Restrictions:

- Do not apply via hand-held equipment.
- DO NOT HANDLE TREATED BRASSICA CROPS for 9 days post-treatment.
- A minimum interval of 7 days must be observed between applications to potato, carrot, parsnip, celeriac, swede, turnip broccoli/calabrese or cauliflower.
- A minimum interval of 10 days must be observed between applications to field bean (harvested dry), beans without pods (dry), combining pea (harvested dry), vining pea (harvested fresh), brussels sprout or cabbage.
- A minimum interval of 7 days must be observed before harvest after applications to vining pea.
- A minimum interval of 28 days must be observed before harvest after applications to field beans.
- A minimum interval of 18 days must be observed before harvest after applications to combining pea.

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE.  
FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**

UFI: KA22-A3GW-X002-20SR

## DIRECTIONS FOR USE

**IMPORTANT:** This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Belanty is a protectant fungicide for the control of a range of diseases in maize, sweetcorn, potatoes, combining pea, vining pea, field bean, beans without pods (dry), cabbage, brussels sprout, broccoli/calabrese, cauliflower, carrot, parsnip, celeriac, swede and turnip.

### Restrictions/Warnings

Belanty has not been fully evaluated on the beneficial organisms used in Integrated Pest Management systems.

Consult processor before using on legumes for processing.

### Disease control

Belanty is a protectant fungicide for the control of a range of diseases in maize, sweetcorn, potatoes, combining pea, vining pea, field bean, beans without pods (dry), cabbage, Brussels sprout, broccoli/calabrese, cauliflower, carrot, parsnip, celeriac, swede and turnip. See summary below:

Apply Belanty as a protectant treatment.

Crops	Targets	Common name	Control
Maize, sweetcorn	<i>Setosphaeria turcica</i>	Leaf blight	C
	<i>Kabatiella zeae</i>	Eyespot	C
Potatoes	<i>Alternaria spp</i>	Alternaria	C
Field bean, beans without pods (dry)	<i>Ascochyta fabae</i>	Ascochyta leaf and pod spot	MC
	<i>Botrytis cinerea, Botrytis fabae</i>	Chocolate spot	MC
	<i>Uromyces viciae fabae</i>	Rust	C
	<i>Ascochyta pisi</i>	Ascochyta leaf and pod spot	MC
Combining pea, vining pea	<i>Botrytis cinerea</i>	Grey mould	MC
	<i>Erysiphe pisi</i>	Powdery mildew	MC

Crops	Targets	Common name	Control
Carrot, parsnip, celeriac	<i>Alternaria dauci</i>	Carrot blight	MC
	<i>Alternaria sp</i>	Alternaria Leaf Blight	MC
	<i>Erysiphe heraclei</i>	Powdery mildew	C
Cabbage	<i>Alternaria sp.</i>	Alternaria	C
	<i>Neopseudocercosporaella brassicae</i>	Ring Spot	C
	<i>Erysiphe sp.</i>	Powdery mildew	C
Broccoli/ calabrese, brussels sprout, cauliflower, swede, turnip	<i>Alternaria sp.</i>	Alternaria	C

C = control

MC = moderate control

### Resistance Management

GROUP	3	FUNGICIDE

Belanty contains mefenitrifluconazole, a DMI fungicide belonging to FRAC Code G1.

Belanty should be used in accordance with FRAC advice for DMIs.

Effective disease management is a critical component in delaying the build-up of resistant pathogen populations. Therefore, Belanty should be applied according to the manufacturers recommended dose and spray timings even when disease pressure is low.

DMIs should be used in mixture, or in alternation with a non-cross resistant fungicide, at a dose rate which is effective against the target disease.

Belanty should be used preventatively and should not be relied upon for its curative potential.

For further advice on resistance management contact your agronomist or specialist adviser or visit the FRAG-UK website.

### **Maize and Sweetcorn**

#### ***Northern Leaf blight (*Setosphaeria turcica*), Eyespot (*Kabatiella zeae*)***

Apply Belanty before the start of disease attack or at early signs of infection.

Apply Belanty at a rate of 1.25 l/ha, in a water volume of 200-300 litres per hectare.

### **Potato**

#### ***Alternaria spp.***

Apply from first basal side shoot visible.

Apply Belanty as a preventative treatment and before disease becomes established in the crop. A maximum of three applications of Belanty may be applied per year with a minimum of 7 days between applications. A minimum of 3 days must elapse between the last spray and harvest.

Apply Belanty at a rate of 1.25 l/ha in a water volume of 200-400 litres per hectare.

### **Field bean, beans without pods (dry), combining pea, vining pea**

Apply Belanty as a preventative treatment and before disease becomes established in the crop or at first signs of disease. A maximum of two applications of Belanty may be applied per year with a minimum of 10 days between applications.

Latest application timing is BBCH 72.

Apply Belanty at a rate of 1.3 L/ha in 100-400 Litres per hectare.

### **Carrot, parsnip and celeriac**

Apply Belanty as a preventative treatment and before disease becomes established in the crop or at first signs of disease. A maximum of three applications of Belanty may be applied per year with a minimum of 7 days between applications. A minimum of 3 days must elapse between the last spray and harvest.

Apply Belanty at a rate of 1.5 l/ha, in a water volume of 200-1000 litres per hectare.

### **Cabbage**

Apply Belanty as a preventative treatment and before disease becomes established in the crop or at first signs of disease. A maximum of three applications of Belanty may be applied per year with a minimum of 10 days between applications. A minimum of 14 days must elapse between the last spray and harvest.

Apply Belanty at a rate of 1.5 l/ha, in a water volume of 200-1000 litres per hectare.

## **Brussels sprouts**

A maximum of three applications of Belanty may be applied per year with a minimum of 10 days between applications. A minimum of 14 days must elapse between the last spray and harvest.

Apply Belanty at a rate of 1.5 l/ha, in a water volume of 200-1000 litres per hectare.

## **Cauliflower, Broccoli/calabrese**

A maximum of three applications of Belanty may be applied per year with a minimum of 7 days between applications. A minimum of 14 days must elapse between the last spray and harvest.

Apply Belanty at a rate of 1.5 l/ha, in a water volume of 200-1000 litres per hectare.

## **Swede and turnip**

A maximum of three applications of Belanty may be applied per year with a minimum of 7 days between applications. A minimum of 3 days must elapse between the last spray and harvest.

Apply Belanty at a rate of 1.5 l/ha, in a water volume of 200-1000 litres per hectare.

## **Following Crops**

Leafy vegetables and fresh herbs, edible flowers, chicory (witloof), herbal infusions from flowers and herbal infusions from leaves and herbs can only be planted/sown 120 days following the application of Belanty. All other crops can be planted/sown following crop failure or following normal commercial harvest of the crops treated with Belanty.

## **Mixing and Application**

### **Mixing**

Never prepare more spray solution than is required.

Three quarters fill the tank with clean water and start the agitation. To ensure thorough mixing of the product, invert the container several times before opening. Add the required quantity of Belanty to the spray tank while re-circulating. Add the remainder of the water and continue agitation until spraying is complete.

When tank mixes are to be used, take due note of any instructions given as to the order of mixing. Each product should be added separately to the spray tank and fully dispersed before the addition of any further product(s).

All tank mixes should be used immediately after mixing.

On emptying the container, rinse container thoroughly by using an integrated pressure rinsing device or by manually rinsing three times. Add washings to tank at time of filling and dispose of container safely.

## **Application**

Apply as a MEDIUM spray, as defined by BCPC.

## **Tank cleaning**

Wash sprayer thoroughly immediately after use, using clean water and following the sprayer cleaning guidance provided by the equipment manufacturer.

## **Compatibility**

For details of compatibilities contact your distributor, local BASF representative or the BASF Technical Services Hotline: 0845 602 2553 or visit our website on: [www.agricentre.bASF.co.uk](http://www.agricentre.bASF.co.uk)

## **The following does not form part of the authorised label text.**

With many products there is a general risk of resistance developing to the active ingredients. For this reason a change in activity cannot be ruled out. It is generally impossible to predict with certainty how resistance may develop because there are so many crop and use connected ways of influencing this. We therefore have to exclude liability for damage or loss attributable to any such resistance that may develop. To help minimise any loss in activity the BASF recommended rate should in all events be adhered to.

Numerous, particularly regional or regionally attributable, factors can influence the activity of the product. Examples include weather and soil conditions, crop plant varieties, crop rotation, treatment times, application amounts, admixture with other products, appearance of organisms resistant to active ingredients and spraying techniques. Under particular conditions a change in activity or damage to plants cannot be ruled out. The manufacturer or supplier is therefore unable to accept any liability in such circumstances. All goods supplied by us are of high grade and we believe them to be suitable, but as we cannot exercise control over their mixing or use or the weather conditions during and after application, which may affect the performance of the material, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use; but nothing should be deemed to exclude or restrict any liability upon us which cannot be excluded or restricted under the provisions of the Unfair Contract Terms Act 1977 or any similar applicable law.

**Additional Product Safety Information**

The product label provides information on a specific pesticidal use of the product; do not use otherwise, unless you have assessed any potential hazard involved, the safety measures required and that the particular use has "off-label" approval or is otherwise permitted under the Plant Protection Product Regulations (EC) No 1107/2009.

The information on this label is based on the best available information including data from test results.

**Great Britain Safety Data Sheet**

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



[agricentre.bASF.co.uk/Belanty/MSDS](http://agricentre.bASF.co.uk/Belanty/MSDS)

Alternatively, contact your supplier.

**Northern Ireland Safety Data Sheet**

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



[agricentre.bASF.co.uk/Belanty-NI/MSDS](http://agricentre.bASF.co.uk/Belanty-NI/MSDS)

Alternatively, contact your supplier.

# Belanty®

**MAPP 20548**

A suspension concentrate containing 75 g/l mefenpropifluconazole. An agricultural/horticultural fungicide for the control of a range of diseases in maize, sweetcorn, potatoes, combining pea, vining pea, field bean, beans without pods (dry), cabbage, Brussels sprout, broccoli/calabrese, cauliflower, carrot, parsnip, celeriac, swede and turnip.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

**SAFETY PRECAUTIONS****Operator protection**

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

DO NOT HANDLE TREATED BRASSICA CROPS for 9 days post-treatment.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH HANDS AND EXPOSED SKIN before meals and after work.

IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

**Environmental protection**

Do not contaminate water with the product or its container.

Do not clean application equipment near surface water.

Avoid contamination via drains from farmyards and roads.

**Storage and disposal**

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place, away from sources of heat.

DO NOT RE-USE CONTAINER FOR ANY PURPOSE. RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times.

Add washings to sprayer at time of filling and dispose of container safely.

Store in a suitable pesticide store. Keep dry and protect from frost.

**This label is compliant with the CPA Voluntary Initiative Guidance****Supplied by:**

BASF plc, 4th and 5th Floors, 2 Stockport Exchange, Railway Road, Stockport, SK1 3GG  
Telephone: 0161 475 3000

Emergency Information (24 hours freephone):  
0049 180 227 3112

Technical Enquiries: 0845 602 2553  
(office hours)

® = Registered trademark of BASF



5 L