

BASF

We create chemistry

Entargo®



MAPP 19312

A suspension concentrate containing 500 g/litre boscalid.

Entargo is a systemic fungicide with protectant properties for disease control in winter wheat, spring wheat, durum wheat, spelt wheat, winter barley and spring barley.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

WHEN USING DO NOT EAT, DRINK OR SMOKE.
WASH HANDS AND EXPOSED SKIN before meals and after work.

Environmental protection

Do not contaminate water with the product or its container.

Do not clean application equipment near surface water.

Avoid contamination via drains from farmyards and roads.

Storage and disposal

KEEP OUT OF REACH OF CHILDREN.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

Do not re-use container for any purpose.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times.

Add washings to sprayer at time of filling and dispose of safely.

Store in a suitable pesticide store. Keep dry and protect from frost.

This label is compliant with the CPA Voluntary Initiative Guidance

5 L

Supplied by:
BASF plc, 4th and 5th Floors, 2 Stockport Exchange, Railway Road,
Stockport SK1 3GG, Telephone: 0161 475 3000

Emergency Information (24 hours freephone): 0049 180 227 3112
Technical Enquiries: 0845 602 2553 (office hours)

© - Registered trademark of BASF

81147438 GB 2100



Entargo®

A suspension concentrate containing 500 g/litre boscalid.

Warning

**May cause an allergic skin reaction.
Toxic to aquatic life with long lasting effects.**

Wear protective gloves/protective clothing/eye protection/face protection.

If medical advice is needed, have product container or label at hand.

Avoid breathing mist.

IF ON SKIN: Wash with plenty of soap and water.

If skin irritation or rash occurs: Get medical advice/attention.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

Contains 1,2-benzisothiazolin-3-one and 2-methyl-4-isothiazolin-3-one. May produce an allergic reaction

To avoid risks to human health and the environment, comply with the instructions for use.



IMPORTANT INFORMATION

FOR USE ONLY AS A PROFESSIONAL AGRICULTURAL FUNGICIDE, as directed below:

Crops	Maximum individual dose	Maximum total dose	Latest time of application
Wheat (winter) Wheat (spring) Wheat (durum) Wheat (spelt) Barley (winter) Barley (spring)	0.7 litre product/ha	0.7 litre product/ha	Before ear emergence (BBCH 51)

Other specific restrictions:

To reduce the risk of resistance developing in target diseases the total number of foliar applications of products containing SDHI fungicides made to any cereal crop must not exceed two.

Entargo must not be applied before BBCH 30.

Do not apply by hand-held equipment.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Restrictions/Warnings

When applied to barley intended for malting, refer to the latest timing restrictions for boscalid given by the British Beer and Pub Association.

Effects of baking have not been fully tested. Consult the grain merchant or processor before use.

You must not apply, more than two foliar applications of products containing complex II inhibitors (SDHIs) to any cereal crop.

After spraying, wash out sprayer thoroughly according to manufacturer's guideline and dispose of washings and clean containers according to DEFRA Code of Practice and local authority guidelines.

Disease control

Disease Control

Entargo is a systemic fungicide with protectant properties for disease control in winter wheat, spring wheat, durum wheat, spelt wheat, winter barley and spring barley as summarised below:

	Winter Wheat	Spring Wheat	Durum Wheat	Spelt Wheat	Winter Barley	Spring Barley
Eyespot	C	C	C	C		
<i>Septoria tritici</i>	MC	MC	MC	MC		
Net blotch					R	R

C = Control

MC = Moderate control

R = Reduction

Resistance Management

Entargo should be used in accordance with the instructions for use for the target diseases at the specific growth stages indicated.

Entargo must always be tank mixed with another fungicide (s) from different cross resistance groups(s)/different mode(s) of action, which when used alone, gives robust control of the same target diseases

Use Entargo as part of an Integrated Crop Management (ICM) strategy incorporating other methods of control.

You must not apply, more than two foliar applications of products containing complex II inhibitors (SDHIs) to any cereal crop.

Entargo should be used preventatively and should not be relied upon for its curative potential.

There is a significant risk of widespread resistance occurring in *Septoria tritici* populations in the UK. Failure to follow resistance management action may result in reduced levels of disease control. For latest advice on resistance management contact your specialist agronomist advisor and visit the FRAG-UK website.

Crop Specific Information

Entargo is a systemic fungicide with protectant properties for disease control in winter wheat, spring wheat, durum wheat, spelt wheat, winter barley and spring barley

Time of Application

For best results apply Entargo as a preventative or protectant treatment. For eyespot control apply when ear is at 1 cm above first node (BBCH 30) up to 2nd node detectable (BBCH 32). For foliar diseases apply before ear emergence (BBCH 51) in wheat and barley.

Rate of Application

Apply 0.7 litre Entargo in a minimum of 100 litres of water per hectare.

Mixing and Application

Mixing

Half fill the spray tank with clean water and start the agitation. SHAKE THE CONTAINER WELL before use and pour in the required amount of product. Rinse any empty containers thoroughly and add rinsings to the spray tank. Add the remainder of the water and continue agitation until spraying is completed.

When tank mixes are to be used, each product should be added separately to the spray tank, taking due note of any instructions given as to the order of mixing.

All tank mixes should be used immediately after mixing.

Application

Apply as a **MEDIUM** spray, as defined by BCPC.

Compatibility

For details of compatibilities contact your distributor, local BASF representative or the BASF Technical Services Hotline: 0845 602 2553 or visit our website on: www.agriCentre.basf.co.uk

The following does not form part of the authorised label text.

With many products there is a general risk of resistance developing to the active ingredients. For this reason a change in activity cannot be ruled out. It is generally impossible to predict with certainty how resistance may develop because there are so many crop and use connected ways of influencing this. We therefore have to exclude liability for damage or loss attributable to any such resistance that may develop. To help minimise any loss in activity the BASF recommended rate should in all events be adhered to.

Numerous, particularly regional or regionally attributable, factors can influence the activity of the product. Examples include weather and soil conditions, crop plant varieties, crop rotation, treatment times, application amounts, admixture with other products, appearance of organisms resistant to active ingredients and spraying techniques. Under particular conditions a change in activity or damage to plants cannot be ruled out. The manufacturer or supplier is therefore unable to accept any liability in such circumstances. All goods supplied by us are of high grade and we believe them to be suitable, but as we cannot exercise control over their mixing or use or the weather conditions during and after application, which may affect the performance of the material, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use; but nothing should be deemed to exclude or restrict any liability upon us which cannot be excluded or restricted under the provisions of the Unfair Contract Terms Act 1977 or any similar applicable law.

Section 6 of the Health and Safety at Work Act

Additional Product Safety Information

The product label provides information on a specific pesticidal use of the product; do not use otherwise, unless you have assessed any potential hazard involved, the safety

measures required and that the particular use has “off-label” approval or is otherwise permitted under the Plant Protection Product Regulations (EC) No 1107/2009.

The information on this label is based on the best available information including data from test results.

Safety Data Sheet

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



agricentre.basf.co.uk/Entargo/MSDS

Alternatively, contact your supplier.

BASF

We create chemistry

Entargo®



MAPP 19312

A suspension concentrate containing 500 g/litre boscalid.

Entargo is a systemic fungicide with protectant properties for disease control in winter wheat, spring wheat, durum wheat, spelt wheat, winter barley and spring barley.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).
WHEN USING DO NOT EAT, DRINK OR SMOKE.
WASH HANDS AND EXPOSED SKIN before meals and after work.

Environmental protection

Do not contaminate water with the product or its container.

Do not clean application equipment near surface water.

Avoid contamination via drains from farmyards and roads.

Storage and disposal

KEEP OUT OF REACH OF CHILDREN.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

Do not re-use container for any purpose.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times.

Add washings to sprayer at time of filling and dispose of safely.

Store in a suitable pesticide store. Keep dry and protect from frost.

This label is compliant with the CPA Voluntary Initiative Guidance

5 L

Supplied by:

BASF plc, 4th and 5th Floors, 2 Stockport Exchange, Railway Road, Stockport SK1 3GG, Telephone: 0161 475 3000

Emergency Information (24 hours freephone): 0049 180 227 3112

Technical Enquiries: 0845 602 2553 (office hours)

© - Registered trademark of BASF

81147438 GB 2100




The
Voluntary
Initiative