

BASF
We create chemistry

Springbok®



MAPP 16786

An emulsifiable concentrate containing 200 g/l metazachlor and 200 g/l dimethenamid-p, for use as a residual herbicide for the control of a range of broad leaved and grass weeds in winter oilseed rape.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) when applying the product.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

WASH CONCENTRATE from skin or eyes immediately.

WASH HANDS before meals and after work.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

IF YOU FEEL UNWELL seek medical advice (show the label if possible).

Do not apply using hand held equipment.

Environmental protection

Do not contaminate surface waters or ditches with chemical or used container.

To protect aquatic organisms respect an unsprayed buffer zone to surface water bodies in line with LERAP requirements.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 metres of the top of the bank of a static or flowing water body, unless a Local Environmental Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 metre from the top of a ditch which is dry at the time of application.

Aim spray away from water.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme.

Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained.

The results of the LERAP must be recorded and kept available for three years.

Storage and disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS. KEEP OUT OF REACH OF CHILDREN.

DO NOT RE-USE CONTAINER for any other purpose.

STORE IN ORIGINAL CONTAINER tightly closed, in a safe place.

On emptying the container, RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times.

Add washings to sprayer at time of filling and dispose of container safely.

This label is compliant with the CPA Voluntary Initiative Guidance

10 L

Supplied by: BASF plc, Agricultural Solutions
PO Box 4, Earl Road, Cheadle Hulme, CHEADLE,
Cheshire SK8 6QG, Tel: 0161 485 6222

Emergency Information: (24 hours freephone): 0049 180 2273112,
Technical Enquiries: 0845 602 2553 (office hours)



The
Voluntary
Initiative

© = Registered trademark of BASF

81132476 GB 1029



Springbok®

An emulsifiable concentrate containing 200 g/l metazachlor and 200 g/l dimethenamid-p,

Danger

Harmful if swallowed.

May be fatal if swallowed and enters airways.

May cause an allergic skin reaction.

Causes serious eye irritation.

Harmful if inhaled.

Suspected of causing cancer.

Very toxic to aquatic life.

Wear protective gloves/clothing/eye protection.

Avoid breathing vapours/spray.

IF SWALLOWED: Immediately call a POISON CENTRE or doctor/physician.

Do NOT induce vomiting.

Store locked up.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.



IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

CROPS:	Winter oilseed rape
MAXIMUM INDIVIDUAL DOSE	2.5 l product/ha
MAXIMUM TOTAL DOSE:	2.5 l product/ha
LATEST TIME OF APPLICATION:	Before 7th true leaf stage (GS 17)
AQUATIC BUFFER ZONE:	5 metres

Other specific restrictions:

Applications shall be limited to a total dose of not more than 1.0kg metazachlor/ha in a three-year period on the same field.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

COMPANY ADVISORY INFORMATION

This section is not part of the Product Label under Regulation (EC) No 1107/2009 and provides additional advice on product use.



Water protection:

This product contains **metazachlor** plus dimethenamid-P and is therefore included in the “**OSR Herbicides? Think Water**” stewardship campaign. Particular care is needed when using in **Surface Water Drinking Water Safeguard Zones**:-

- **Do not use** after 30 September on drained fields, including mole-drained.
- In other areas:-
- Avoid use after 15 October on drained fields, including mole-drained.

For further information, see www.agriculture.basf.co.uk and www.osrherbicides.org.uk or telephone BASF on 0845 602 2553.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

SPRINGBOK is a residual herbicide that can be used on all varieties of winter oilseed rape for the control of a range of broad leaved and grass weeds.

Restrictions/Warnings

When SPRINGBOK is applied pre emergence, it is important for crop safety to ensure physical separation of the herbicide from the seed. Ensure that the seed is well covered with soil to a depth of 15 mm. With direct drilled crops, harrow across the slits to cover the seed before spraying.

Do not disturb the soil after application.

Do not apply to broadcast crops until the crop has attained the two fully expanded cotyledon growth stage.

Do not use SPRINGBOK on sands, very light soils or soils containing more than 10% organic matter.

Do not apply to stony soils; i.e. stones, flints or chalk readily visible on surface. On brashy and stony soils, SPRINGBOK may cause some reduction in crop vigour and/or plant stand.

Do not apply to cloddy seedbeds. Seedbeds must have a fine, firm tilth for optimum weed control. Loose or cloddy seedbeds must be consolidated prior to application.

Do not apply when heavy rain is forecast and do not use on waterlogged soil or soils prone to waterlogging. Crop thinning or reductions in crop vigour can occur if there is very wet weather after application.

Soil moisture is required for effective weed control via root uptake. Residual control may be reduced under prolonged dry conditions.

Weeds germinating from depth may not be controlled.

Do not apply SPRINGBOK to crops suffering from stress, which may be caused, for example, by pests, disease, waterlogging, poor seedbed conditions or previous chemical treatment.

Under frosty conditions a transient scorch may occur.

Extreme care should be taken to avoid damage by drift onto plants outside the target area.

To reduce the risk of movement of Springbok to water:

- a. On clay soils, create a fine, consolidated seedbed to slow the downward movement of water.
- b. Do not apply Springbok to dry soil. Moist soils have fewer and smaller cracks.
- c. Do not apply Springbok if heavy rain is forecast, wait until after the event.

Weed control

SPRINGBOK is taken up via cotyledons and roots and takes maximum effect before, or shortly after, weed emergence. Optimum results are obtained from applications made to fine, firm and moist seedbeds.

Susceptibility of weeds to single applications of SPRINGBOK.

Weed species	SPRINGBOK 2.5 l/ha	
	Application pre-emergence of weed. Weed susceptibility rating	Application post-emergence of weed. Maximum susceptible growth stage of weed
Black-grass	MR	Pre-em
Chickweed, Common	S	4 true leaf
Cleavers	S	Pre-em
Crane's-bill, Cut-leaved	S	Cotyledon
Crane's-bill, Round-leaved	S	Cotyledon
Crane's-bill, Small-flowered	S	Cotyledon
Forget-me-not, Field	MS	Pre-em
Mayweed, Scented	S	Pre-em
Pansy, Field	MR	Pre-em
Poppy, Common	S	Pre-em
Shepherd's-purse	S	1 true leaf (MS)
Speedwell, Common Field	S	1 true leaf

S = Susceptible

MS = Moderately Susceptible

MR = Moderately Resistant

R = Resistant

Pre-em = Pre-emergence of weed

Resistance Management

Strains of some annual grasses (e.g. black-grass, wild-oats, and Italian rye-grass) have developed resistance to herbicides which may lead to poor control. A strategy for preventing and managing such resistance should be adopted. Guidelines have been produced by the Weed Resistance Action

Group (WRAG) and copies are available from the HGCA, CPA, your distributor, crop adviser or product manufacturer.

Repeated use of herbicides with the same mode of action can increase the risk of strains of weeds developing resistance to these compounds, leading to poor control. In order to minimise the risk, a strategy for preventing and managing such resistance should be adopted. Use products with different modes of action and from different chemical groups in sequence, in conjunction with effective cropping rotation and cultivation techniques. In all cases the recommended rate of use should be adhered to. Follow WRAG guidelines.

Key elements of the resistance management strategy for SPRINGBOK.

- Always follow WRAG guidelines for preventing and managing herbicide resistant weeds.
- Maximise the use of cultural control measures wherever possible (e.g. crop rotation, ploughing, stale seedbeds, delayed drilling, etc).
- Use sequences of effective herbicides with different modes of action within individual crops, or successive crops.
- For the control of black-grass, always use SPRINGBOK in sequence with other effective graminicides with different modes of action.
- Monitor fields regularly and investigate the reasons for any poor control.

Crop Specific Information

SPRINGBOK can be used on all varieties of winter oilseed rape.

Time of application

SPRINGBOK may be applied pre or post emergence of the crop but for optimum activity application must be pre emergence of the weeds.

Extreme care must be taken to avoid spray drift onto non-crop plants outside the target area.

Pre-crop emergence applications should only be made on medium and heavy soils (as defined by the ADAS Soil Texture (85) System). Pre-emergence applications should be made before the crop seed chits, which in optimum conditions may occur within 48 hours of drilling.

DO NOT USE SPRINGBOK pre-crop emergence when any of the following conditions apply:

- When crop has been broadcast.
- The crop seed has started to germinate (i.e. 48 hours after drilling)
- Seed is not covered with 15 mm of soil;
- When heavy rain is forecast;
- Dry, cloddy or open seedbeds;
- Late drilled crops – as a guide these are crops drilled later than the first week in September in Northern England and Scotland and later than mid-September in the rest of the UK;
- Stony soils, i.e. stones, flints or chalk are readily visible on the soil surface.

Post-crop emergence applications can be made in the following circumstances:

- When the majority of the crop has two fully expanded cotyledons and up to before the seventh true leaf growth stage (GS 17);
- On light, medium and heavy soils (as defined by the ADAS Soil Texture (85) System);
- When seedbed conditions are not ideal for pre-emergence use;

- When the seed is not covered with 15 mm of soil (as with broadcast crops);
- To light, stony, brashy or gravelly soils, particularly if heavy rainfall was forecast for the period following drilling;
- When late-drilled crops are to be treated.

Rate of application

Apply 2.5 litres/hectare SPRINGBOK in 200-300 litres water volume per hectare.

Extension of Authorisation

This extension of the authorised use (number 1540 of 2015) provides for the use of Springbok in respect of crops and situations other than those included on the product label (above). Neither the efficacy or the phytotoxicity of the product for which this Extension of authorisation has been granted have been assessed and, as such, the user bears the risk in respect of failures concerning its efficacy and phytotoxicity.

The following Extension of Use is authorised for Springbok applied post drilling but pre-crop emergence or up to the latest time of application for transplanted crops.

Crop(s)	Max. individual dose (l/ha)	Max. total dose (l/ha)	Latest time of application	Buffer zone
Outdoor cabbage, cauliflower, broccoli/calabrese and Brussels sprouts	2.5	2.5	Up to 7 days post transplanting	5 metres
Outdoor crops of kale, collard, chinese leaf cabbage, oriental cabbages. tat soi, pak choi, choi sum	2.5	2.5	10 days after transplanting or before 8 true leaves for drilled crops	5 metres
Outdoor kohlrabi	2.5	2.5	Before 9 true leaves unfolded	5 metres

Advisory Information

In field brassicas, optimum results have been obtained pre-emergence of the weed.

Pre-emergence applications of Springbok to direct drilled brassicas may cause damage especially at higher doses.

HDC trials in 2005 and 2006 demonstrated the crop safety of Springbok to cauliflower when applied at 2.5 l/ha 1-2 days post-transplanting. However, this 'extension of use' remains at the risk of the user.

Springbok should be applied in a minimum water volume of 220 litres water/ha.

Irrigation soon after application may result in crop damage.

A minimum PHI of 32 days must be observed.

Following Crops

Any crop can follow normally harvested winter oilseed rape treated with SPRINGBOK. Ploughing is not essential before sowing a following cereal crop, but is required for all other crops.

In the event of crop failure, the following crops may be sown after ploughing to a depth of at least 15 cm:

Redrilling the same autumn:	winter wheat (excluding durum) or winter barley;
Redrilling the following spring:	cereals (excluding durum wheat), spring oilseed rape, peas and field beans.

Mixing and Application

Application

SPRINGBOK should be applied in a water volume of 200-300 litres per hectare
Apply as a **MEDIUM** spray, as defined by BCPC.

Mixing

Never prepare more spray solution than is required.

SHAKE WELL BEFORE USE.

Fill the spray tank three quarters full with water and start the agitation.

To ensure thorough mixing of the product, invert the container several times before opening.

Add the required quantity of SPRINGBOK to the spray tank while re-circulating.

Fill up the tank with water and continue agitation until spraying is completed.

On emptying the product container, rinse container thoroughly by using an integrated pressure rinsing device or manually rinsing three times.

Add washings to sprayer at time of filling and dispose of container safely.

Tank cleaning

Thoroughly rinse application equipment three times with water after use.

The following does not form part of the authorised label text.

With many products there is a general risk of resistance developing to the active ingredients. For this reason a change in activity cannot be ruled out. It is generally impossible to predict with certainty how resistance may develop because there are so many crop and use connected ways of influencing this. We therefore have to exclude liability for damage or loss attributable to any such resistance that may develop. To help minimise any loss in activity the BASF recommended rate should in all events be adhered to.

Numerous, particularly regional or regionally attributable, factors can influence the activity of the product. Examples include weather and soil conditions, crop plant varieties, crop rotation, treatment times, application amounts, admixture with other products, appearance of organisms resistant to active ingredients and spraying techniques. Under particular conditions a change in activity or damage to plants cannot be ruled out. The manufacturer or supplier is therefore unable to accept any liability in such circumstances. All goods supplied by us are of high grade and we believe them to be suitable, but as we cannot exercise control over their mixing or use or the weather conditions during and after application, which may affect the performance of the material, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use; but nothing should be deemed to exclude or restrict any liability upon us which cannot be excluded or restricted under the provisions of the Unfair Contract Terms Act 1977 or any similar applicable law.

Section 6 of the Health and Safety at Work Act

Additional Product Safety Information

The product label provides information on a specific pesticidal use of the product; do not use otherwise, unless you have assessed any potential hazard involved, the safety measures required and that the particular use has “off-label” approval or is otherwise permitted under the Plant Protection Products Regulations.

The information on this label is based on the best available information including data from test results.

Safety Data Sheet

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



bit.ly/Springbok_sds

Alternatively, contact your supplier.

BASF
We create chemistry

Springbok®



MAPP 16786

An emulsifiable concentrate containing 200 g/l metazachlor and 200 g/l dimethenamid-p, for use as a residual herbicide for the control of a range of broad leaved and grass weeds in winter oilseed rape.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) when applying the product.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

WASH CONCENTRATE from skin or eyes immediately.

WASH HANDS before meals and after work.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

IF YOU FEEL UNWELL seek medical advice (show the label if possible).

Do not apply using hand held equipment.

Environmental protection

Do not contaminate surface waters or ditches with chemical or used container.

To protect aquatic organisms respect an unsprayed buffer zone to surface water bodies in line with LERAP requirements.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 metres of the top of the bank of a static or flowing water body, unless a Local Environmental Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 metre from the top of a ditch which is dry at the time of application.

Aim spray away from water.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme.

Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained.

The results of the LERAP must be recorded and kept available for three years.

Storage and disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS. KEEP OUT OF REACH OF CHILDREN.

DO NOT RE-USE CONTAINER for any other purpose.

STORE IN ORIGINAL CONTAINER tightly closed, in a safe place.

On emptying the container, RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times.

Add washings to sprayer at time of filling and dispose of container safely.

This label is compliant with the CPA Voluntary Initiative Guidance

10 L

Supplied by: BASF plc, Agricultural Solutions
PO Box 4, Earl Road, Cheadle Hulme, CHEADLE,
Cheshire SK8 6QG, Tel: 0161 485 6222

Emergency Information: (24 hours freephone): 0049 180 2273112,
Technical Enquiries: 0845 602 2553 (office hours)



The
Voluntary
Initiative

© = Registered trademark of BASF

81132476 GB 1029

