



Divexo[®]



MAPP 17937

**A suspension concentrate containing 375 g/litre chlorothalonil and 37.5 g/l fluxapyroxad
A protectant fungicide for use in winter wheat, spring wheat, durum wheat, spelt wheat, winter barley and spring barley**

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

AVOID ALL CONTACT WITH SKIN.

WASH HANDS AND EXPOSED SKIN before meals and after work.

WHEN USING DO NOT EAT, DRINK OR SMOKE. IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

Environmental protection

Do not contaminate water with the product or its container. Do not clean equipment near surface water. Avoid contamination via drains from farmyards and roads.

To protect aquatic organisms respect an unsprayed buffer zone distance to surface water bodies in line with LERAP requirements. **DO NOT ALLOW DIRECT SPRAY** from horizontal boom sprayers to fall within 5m of the top of the bank of a static or flowing waterbody, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1m of the top of a ditch which is dry at the time of application. Aim spray away from water.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom

sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for inspection for three years.

Storage and disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

Store in a suitable pesticide store. Keep dry and protect from frost.

10 L e

This label is compliant with the CPA Voluntary Initiative Guidance

Supplied by:
BASF plc
Crop Protection
PO Box 4, Earl Road
Cheadle Hulme, CHEADLE
Cheshire SK8 6QG
Tel: 0161 485 6222
Emergency Information:
(24 hours freephone):
0049 180 2273112
Technical Enquiries:
0845 602 2553 (office hours)

 **BASF**
We create chemistry

© = Registered trademark of BASF



The
Voluntary
Initiative



81121575GB1107

Divexo®

A suspension concentrate containing
375 g/litre chlorothalonil and 37.5 g/l fluxapyroxad

Warning:

Harmful if inhaled.
May cause respiratory irritation.
May cause an allergic skin reaction.
Suspected of causing cancer.
Very toxic to aquatic life with long lasting effects.

Wear suitable protective clothing (coveralls) and suitable protective gloves when handling the concentrate.
Use only outdoors or in a well-ventilated area.
Obtain special instructions before use.

IF exposed or concerned: Call a POISON CENTRE or doctor/physician.

Store in a well-ventilated place. Keep container tightly closed.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.



To avoid risks to human health and the environment, comply with the instructions for use.

IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE, as directed below:

Crops	Maximum individual dose	Maximum total dose	Latest time of application	Aquatic buffer zone distance
Winter wheat, spring wheat, durum wheat, spelt wheat, winter barley and spring barley	2.0 litres product/ha	2.0 litres product/ha	Up to and including emergence of ear just complete (GS 59)	5 metres

Other specific restriction:

This product must not be applied via hand-held equipment.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Divexo is a fungicide with protectant properties for disease control in winter wheat, spring wheat, durum wheat, spelt wheat, winter and spring barley.

1. Restrictions/Warnings

No more than two foliar applications of any SDHI fungicide may be made to any cereal crop.

When applied to barley intended for malting, refer to the latest timing restrictions for fluapyroxad given by the British Beer and Pub Association.

2. Disease Control

Divexo is a protectant fungicide for disease control in winter wheat, spring wheat, durum wheat, spelt wheat, winter and spring barley as summarised below:

	Winter Wheat	Spring Wheat	Durum Wheat	Spelt Wheat	Winter Barley	Spring Barley
<i>Septoria tritici</i>	C	C	C	C	-	-
Net blotch	-	-	-	-	C	C
Ramularia	-	-	-	-	C	C
<i>Rhynchosporium</i>	-	-	-	-	C	C

C = Control

Note:

Apply Divexo as a preventative or protectant treatment, do not apply to established infections or rely on any curative potential

2.1 Resistance Management

Divexo contains fluxapyroxad a member of the SDHI cross resistance group. Use Divexo as part of an Integrated Crop Management (ICM) strategy incorporating other methods of control, including where appropriate other fungicides with a different mode of action. You must not apply more than two foliar applications of SDHI containing products to any cereal crop. For further advice on resistance management in SDHIs contact your agronomist or specialist advisor, and visit the FRAG-UK website.

3. Crop specific information

3.1 Crops

Divexo is a fungicide with protectant properties for disease control in winter wheat, spring wheat, durum wheat, spelt wheat, winter and spring barley.

3.2 Time of Application

For best results apply Divexo as a preventative treatment or at first sign of foliar disease attack. A maximum of one application can be made up to and including emergence of ear just complete (GS 59) in wheat and barley.

3.3 Rate of Application

Apply 2.0 litres Divexo in a minimum of 200 litres of water per hectare.

3.4 Qualified recommendation

Divexo may be applied at 2 litres per hectare in 100 litres water per hectare although efficacy at this reduced volume has not been evaluated. Therefore application at 100 litres water per hectare is at user's risk with regard to biological efficacy.

4. Mixing and Spraying

Apply as a MEDIUM spray, as defined by BCPC.

Half fill the spray tank with clean water and start the agitation. SHAKE THE CONTAINER WELL before use and pour in the required amount of product. Rinse any empty containers thoroughly and add rinsings to the spray tank. Add the remainder of the water and continue agitation until spraying is completed.

After spraying, wash out sprayer thoroughly according to manufacturer's guideline and dispose of washings and clean container according to DEFRA Code of Practice and local authority guidelines

When tank mixes are to be used, each product should be added separately to the spray tank, taking due note of any instructions given as to the order of mixing.

All tank mixes should be used immediately after mixing.

5. Compatibility

For details of compatibilities contact your distributor, local BASF representative or the BASF Technical Services Hotline: 0845 602 2553 or visit our website on: www.agriCentre.basf.co.uk.

6. Trademarks

Divexo® is a registered trademark of BASF.

SPECIMEN

The following does not form part of the product label under the Plant Protection Product Regulations (EC) No 1107/2009:-

With many products there is a general risk of resistance developing to the active ingredients. For this reason a change in activity cannot be ruled out. It is generally impossible to predict with certainty how resistance may develop because there are so many crop and use connected ways of influencing this. We therefore have to exclude liability for damage or loss attributable to any such resistance that may develop. To help minimise any loss in activity the BASF recommended rate should in all events be adhered to.

Numerous, particularly regional or regionally attributable, factors can influence the activity of the product. Examples include weather and soil conditions, crop plant varieties, crop rotation, treatment times, application amounts, admixture with other products, appearance of organisms resistant to active ingredients and spraying techniques. Under particular conditions a change in activity or damage to plants cannot be ruled out. The manufacturer or supplier is therefore unable to accept any liability in such circumstances. All goods supplied by us are of high grade and we believe them to be suitable, but as we cannot exercise control over their mixing or use or the weather conditions during and after application, which may affect the performance of the material, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use; but nothing should be deemed to exclude or restrict any liability upon us which cannot be excluded or restricted under the provisions of the Unfair Contract Terms Act 1977 or any similar applicable law.

Section 6 of the Health and Safety at Work Act Additional Product Safety Information

The product label provides information on a specific pesticidal use of the product; do not use otherwise, unless you have assessed any potential hazard involved, the safety measures required and that the particular use has “off-label” approval or is otherwise permitted under the Plant Protection Products Regulation (EC) No 1107/2009.

The information on this label is based on the best available information including data from test results

Safety Data Sheet

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



bit.ly/Divexo_sds

Alternatively, contact your supplier.