

PicoPro®



MAPP 13454

A suspension concentrate containing 16 g/l picolinafen and 320 g/l pendimethalin.

A herbicide with residual and contact activity for the control of meadow grasses and broad leaved weeds in winter and spring wheat and winter and spring barley, rye and triticale.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves

WASH HANDS before eating and after work

IF YOU FEEL UNWELL seek medical advice (show the label if possible)

Environmental protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

To protect aquatic organisms respect an unsprayed buffer zone to surface water bodies. DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing water body, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1m of the top of a ditch which is dry at the time of application, Aim spray away from water.

This product qualifies for inclusion within the Local Environment Risk Assessment for

Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years.

Storage and disposal

DO NOT RE-USE CONTAINER for any other purpose

On emptying the container, RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times.

Add washings to sprayer at time of filling and dispose of container safely.

STORE IN ORIGINAL CONTAINER tightly closed, in a safe place.

PROTECT FROM FROST

10 Le



The
Voluntary
Initiative

This label is compliant with the CPA
Voluntary Initiative Guidance

Supplied by:
BASF plc
Crop Protection
PO Box 4, Earl Road
Cheadle Hulme, CHEADLE
Cheshire SK8 6QG
Tel: 0161 485 6222
Emergency Information:
(24 hours freephone):
0049 180 2273112
Technical Enquiries:
0845 602 2553 (office hours)

 **BASF**
We create chemistry

© registered trademark of BASF

81117908GB1067



PicoPro®

A suspension concentrate containing 16 g/litre picolinafen,
320 g/litre pendimethalin and 1, 2-benzisothiazole



WARNING:
VERY TOXIC TO AQUATIC LIFE WITH LONG LASTING EFFECTS.

COLLECT SPILLAGE.
DISPOSE OF CONTENTS/CONTAINER TO A LICENSED HAZARDOUS-WASTE DISPOSAL CONTRACTOR OR COLLECTION SITE EXCEPT FOR EMPTY CLEAN CONTAINERS WHICH CAN BE DISPOSED OF AS NON-HAZARDOUS WASTE.

CONTAINS PENDIMETHALIN. CONTAINS 1, 2-BENZISOTHIAZOLE
MAY PRODUCE AN ALLERGIC REACTION

To avoid risks to human health and the environment, comply with the instructions for use.

This product is approved under the Plant Protection Products Regulations 2005.

IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

Crop	Application Timing	Maximum Individual Dose (litres product/ha)	Maximum Number of Treatments (per crop)	Latest Time of Application
Winter barley, winter wheat, rye, triticale.	Pre- and post-emergence	3.0	1	Before pseudo-stem erect stage (before growth stage 30)
Spring barley Spring wheat	Pre-emergence OR Post-emergence	3.0 OR 2.0	1	Pre-emergence of the crop OR Before pseudo-stem erect stage (before growth stage 30)

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

PicoPro is a herbicide with residual and contact activity for the control of meadow grasses and broad leaved weeds in winter and spring wheat and winter and spring barley, rye and triticale.

1. Restrictions/Warnings

1.1 Efficacy

For residual control some soil moisture is required for PicoPro. Best results will be obtained if rain falls within 7 days of application.

Residual control may be reduced :

- under prolonged dry conditions
- on soils with more than 6% organic matter
- on soils with a high Kd factor
- on soils where ash content is high

DO NOT disturb the soil after application.

Where cultural techniques which encourage the build up of organic residues in the surface soil are practised for a number of seasons, the effectiveness of residual herbicides may be reduced. In such circumstances periodic ploughing is recommended to disperse residues into a greater volume of soil.

1.2 Soil types

PicoPro may be used on all mineral soil types.

DO NOT use on soils with more than 10% organic matter.

DO NOT use on stony or gravelly soils as crop damage can occur, particularly if heavy rain follows treatment.

DO NOT use on water logged soil or soils prone to water logging.

1.3 Seedbed preparation

Trash and straw should be incorporated evenly during seedbed preparation.

Seedbeds must have a fine, firm tilth.

Loose or cloddy seedbeds must be consolidated prior to application, otherwise reduced weed control may occur. They may also cause seed to be inadequately covered which could result in crop damage.

1.4 Crop safety

Do not apply to crops suffering from stress, which may be caused for example by pests, disease, waterlogging, poor seedbed conditions or previous chemical treatment.

Water logging that occurs soon after pre-emergence applications may lead to crop damage.

For pre-emergence applications, seed should be covered with a minimum of 3.2cm of settled soil.

Shallow drilled crops should only be treated post-emergence

DO NOT soil incorporate.

Avoid spraying during periods of prolonged or severe frosts.

Some transient bleaching may be seen after application to some crops. This does not lead to yield loss.

DO NOT spray undersown cereals or those to be undersown.

DO NOT roll emerged crops prior to application.

DO NOT roll autumn treated crops until the spring.

1.5 Spray drift

Extreme care should be taken to avoid damage by drift onto plants outside the target area.

1.6 Other restrictions/warnings

DO NOT use more than one application of PicoPro in one growing season.

Concentrated or diluted PicoPro may stain. Avoid spillage. Staining is minimised or completely removed if skin/clothes are washed immediately.

2. Weed Control

2.1 Susceptibility of weeds to single applications of PicoPro

For post-emergence applications best results will be achieved by application to small actively growing weeds.

Product Rate (l/ha)	PicoPro		
	3	2	3
Application timing	Pre-em	Post-em	
Annual Meadow-grass	S	S up to 2 lvs	S up to 2 lvs
Rough Meadow-grass	MS	MS up to 2 lvs	MS up to 2 lvs
Loose silky bent	S	MR up to 1 leaf	S up to 1 leaf
Charlock	MS	S up to 6 lvs	S up to 6 lvs
Cleavers	MS	MS up to 1 whorl	S up to 3 whorls
Common Chickweed	S	S up to 6 lvs	S up to 6 lvs
Common Field Speedwell	S	S up to 6 lvs	S up to 6 lvs
Common Fumitory	S	S up to 3 lvs	S up to 3 lvs
Common Orache	MS	-	-
Common Poppy	MS	S up to 6 lvs	S up to 6 lvs
Corn Marigold	S	-	-
Fat-hen	MS	S up to 2 lvs	S up to 2 lvs
Field Forget-me-not	MS	S up to 1 leaf	S up to 2 lvs
Field Pansy	MS	S up to 4 lvs	S up to 4 lvs
Hemp-nettle (Day Nettle)	S	-	-
Herbit Dead-nettle	S	-	S up to 6 lvs
Ivy-leaved speedwell	S	S up to 6 lvs	S up to 6 lvs
Knotgrass	MS	-	-
Parsley Piert	S	-	S up to 2 lvs
Red Dead-nettle	S	S up to 6 lvs	S up to 6 lvs
Scarlet Pimpernel	S	-	-
Shepherd's Purse	S	S up to 6 lvs	S up to 6 lvs
Smooth Sowthistle	MS	-	-
Volunteer Oilseed Rape (1)	S	MS up to 2 lvs	MS up to 2 lvs

S = Susceptible
MR = Moderately resistant

MS = Moderately susceptible
(1) = deep germinating volunteer Oilseed Rape may not be controlled.

2.2 Resistance management

Strains of some annual grasses (eg Blackgrass, Wild-Oats, and Italian Ryegrass) have developed resistance to herbicides, which may lead to poor control. Picolinafen is an inhibitor of carotenoid biosynthesis at the PDS step (a group that includes picolinafen and diflufenican). A strategy for preventing and managing such resistance should be adopted. To prevent the development of resistant weeds herbicides with different modes of action must be used when applying in sequence. Guidelines have been produced by the Weed Resistance Action Group and copies are available from the AHDB, CPA, your distributor, crop adviser or product manufacturer.

3. Crop specific information

3.1 Crops

All varieties of Winter and Spring Wheat, Winter and Spring Barley, Rye and Triticale.

3.2 Time of application

Pre- and post-crop emergence, up to before pseudo-stem erection (before growth stage 30).

For winter drilled crops DO NOT apply pre-emergence to crops drilled after 30 November.

3.3 Rate of application

Apply PicoPro at 2 or 3 l/ha in 200 litres of water per hectare.

4. Following crops

4.1 Following autumn use of PicoPro

Following crops after normal harvest

There are no restrictions on following crops other than rye-grass when PicoPro is used alone. For ryegrass, ploughing or cultivation to at least 15 cm is required prior to drilling.

In the event of crop failure

Spring wheat, spring barley, and winter field beans can be sown after crop failure. Plough to at least 15 cm and allow a minimum of 8 weeks between using PicoPro and sowing.

Maize, oilseed rape, peas, or spring field beans can be sown in spring after crop failure. Minimal non-inversion soil cultivation (to 5 cm) is required before drilling. At least 5 months must elapse between using PicoPro and sowing maize.

4.2 Following Spring use of PicoPro

Following crops after normal harvest

Oilseed rape, winter field beans, winter cereals, and ryegrass can be sown in autumn following normal harvest. Minimal non-inversion soil cultivation (to 5 cm) is needed before the sowing of oilseed rape or winter field beans. Ploughing to a depth of at least 15 cm is needed before sowing winter cereals and ryegrass. At least 6 months must elapse between using PicoPro and sowing ryegrass.

Maize, spring field beans, peas, spring wheat, and spring barley can be sown in spring following normal harvest. Minimal non-inversion soil cultivation (to 5 cm) is needed before sowing.

In the event of crop failure

Spring wheat, spring barley, peas, spring field beans, maize, and spring oilseed rape can be sown after crop failure. Plough to at least 15 cm and allow a minimum of 8 weeks to elapse between using PicoPro and sowing.

5. Mixing and Spraying

5.1 Mixing

Mixing and Spraying

Apply PicoPro in 200 litres water/ha.

Ensure good, even spray cover of the target using a spray of **fine** or **medium** quality (BCPC definition).

Never prepare more spray solution than is required.

Fill the spray tank three quarters full with water and start the agitation. To ensure thorough mixing of the product, invert the container several times before opening. Add the required quantity of PicoPro to the spray tank while re-circulating and then the remainder of the water. Continue agitation until spraying is completed.

On emptying the product container, rinse container thoroughly by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of container safely.

5.2 Sprayer cleaning

After spraying, thoroughly clean and flush out application machinery with a minimum of three rinses, using a wetting agent or a proprietary spray tank cleaner to ensure that all traces of product are removed, particularly before using the sprayer in broad-leaved crops.

6. Compatibility

Contact your distributor or BASF representative for compatibility information.

7. Trademark acknowledgements

PicoPro is a registered trademark of BASF.

The following does not form part of the product label under the Plant Protection Products Regulations.

With many products there is a general risk of resistance developing to the active ingredients. For this reason a change in activity cannot be ruled out. It is generally impossible to predict with certainty how resistance may develop because there are so many crop and use connected ways of influencing this. We therefore have to exclude liability for damage or loss attributable to any such resistance that may develop. To help minimise any loss in activity the BASF recommended rate should in all events be adhered to.

Numerous, particularly regional or regionally attributable, factors can influence the activity of the product. Examples include weather and soil conditions, crop plant varieties, crop rotation, treatment times, application amounts, admixture with other products, appearance of organisms resistant to active ingredients and spraying techniques. Under particular conditions a change in activity or damage to plants cannot be ruled out. The manufacturer or supplier is therefore unable to accept any liability in such circumstances. All goods supplied by us are of high grade and we believe them to be suitable, but as we cannot exercise control over their mixing or use or the weather conditions during and after application, which may affect the performance of the material, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use; but nothing should be deemed to exclude or restrict any liability upon us which cannot be excluded or restricted under the provisions of the Unfair Contract Terms Act 1977 or any similar applicable law.

**Section 6 of the Health and Safety at Work Act
Additional Product Safety Information**

The product label provides information on a specific pesticidal use of the product; do not use otherwise, unless you have assessed any potential hazard involved, the safety measures required and that the particular use has 'off-label' approval or is otherwise permitted under the Plant Protection Products Regulations 2005.

The information on this label is based on the best available information including data from test results

Safety Data Sheet

To access the Safety Data Sheet for this product scan the QR code or use the weblink below.



bit.ly/PicoPro_sds

Alternatively, contact your supplier.

SPECIMEN

PicoPro®



MAPP 13454

A suspension concentrate containing 16 g/l picolinafen and 320 g/l pendimethalin.

A herbicide with residual and contact activity for the control of meadow grasses and broad leaved weeds in winter and spring wheat and winter and spring barley, rye and triticale.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves

WASH HANDS before eating and after work

IF YOU FEEL UNWELL seek medical advice (show the label if possible)

Environmental protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

To protect aquatic organisms respect an unsprayed buffer zone to surface water bodies. DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing water body, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1m of the top of a ditch which is dry at the time of application, Aim spray away from water.

This product qualifies for inclusion within the Local Environment Risk Assessment for

Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years.

Storage and disposal

DO NOT RE-USE CONTAINER for any other purpose

On emptying the container, RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of container safely.

STORE IN ORIGINAL CONTAINER tightly closed, in a safe place.

PROTECT FROM FROST

10 Le

This label is compliant with the CPA
Voluntary Initiative Guidance



The
Voluntary
Initiative

Supplied by:
BASF plc
Crop Protection
PO Box 4, Earl Road
Cheadle Hulme, CHEADLE
Cheshire SK8 6QG
Tel: 0161 485 6222
Emergency Information:
(24 hours freephone):
0049 180 2273112
Technical Enquiries:
0845 602 2553 (office hours)

 **BASF**
We create chemistry

© registered trademark of BASF

81117908GB1067