

Jenton®



MAPP 11898

An emulsifiable concentrate containing 100 g/litre (10.1% w/w) pyraclostrobin and 375 g/litre (37.9% w/w) fenpropimorph.

A protectant fungicide with curative properties for use in winter wheat, winter barley, spring wheat, spring barley and oats.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective clothing:

WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

AVOID ALL CONTACT WITH SKIN.

WASH HANDS AND EXPOSED SKIN before meals and after work. **WHEN USING DO NOT EAT, DRINK OR SMOKE.**

IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

Environmental protection

Do not contaminate surface waters or ditches with chemical or used container.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing

waterbody, unless a Local Environmental Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application. Aim spray away from water.

This product qualifies for inclusion within the Local Environmental Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years.

Storage and disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to spray at time of filling and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

Store in a suitable pesticide store. Keep dry and protect from frost.



5 Le

UN 3082
Environmentally hazardous
substance, liquid, N.O.S.
(contains pyraclostrobin 10%
and fenpropimorph 38%)
Marine Pollutant

This label is compliant
with the CPA Voluntary
Initiative Guidance



Supplied by:
BASF plc
Crop Protection
PO Box 4, Earl Road
Cheadle Hulme, CHEADLE
Cheshire SK8 6QG
Tel: 0161 485 6222
Emergency Information:
(24 hours freephone):
0049 180 2273112
Technical Enquiries:
0845 602 2553 (office hours)

BASF
We create chemistry

81104210GB1016



® = Registered trademark of BASF




Jenton®

An emulsifiable concentrate containing
100 g/litre (10.1% w/w) pyraclostrobin and
375 g/litre (37.9% w/w) fenpropimorph

DANGER
HARMFUL IF INHALED
HARMFUL IF SWALLOWED
CAUSES SKIN IRRITATION
CAUSES SERIOUS EYE IRRITATION
SUSPECTED OF DAMAGING THE
UNBORN CHILD
MAY BE FATAL IF SWALLOWED AND
ENTERS AIRWAYS
VERY TOXIC TO AQUATIC LIFE
WITH LONG LASTING EFFECTS

OBTAIN SPECIAL INSTRUCTIONS BEFORE USE
AVOID BREATHING MIST
WEAR PROTECTIVE GLOVES AND EYE/FACE PROTECTION
IF SWALLOWED: IMMEDIATELY CALL A POISON CENTER OR
DOCTOR/PHYSICIAN
DO NOT INDUCE VOMITING
IF ON SKIN: WASH WITH PLENTY OF SOAP AND WATER

**To avoid risks to human health and the environment, comply
with the instructions for use.**



This product is approved under the Control Of Pesticide Regulations 1986.

IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE, as directed below:

Crops	Maximum individual dose	Maximum number of treatments	Latest time of application
Winter and spring wheat	2.0 litres product per hectare	2 per crop	Up to and including flowering (anthesis) complete (GS 69)
Winter and spring barley Oats	2.0 litres product per hectare	2 per crop	Up to and including emergence of ear just complete (GS 59)

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.



DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product. Jenton is a fungicide with protectant and curative properties for use in winter wheat, winter barley, spring wheat, spring barley and oats for disease control. Yield response may be obtained in the absence of visual disease.

1. Restrictions/Warnings

Extreme care must be taken to avoid drift onto nearby crops. Wash equipment thoroughly after use.

2. Disease Control

Jenton is a fungicide with protectant and curative properties for use in winter wheat, winter barley, spring wheat, spring barley and oats for disease control as summarised below:

	Winter Wheat	Spring Wheat	Winter Barley	Spring Barley	Oats
<i>Septoria tritici</i>	C	C			
Mildew			C	C	C
Yellow rust	C	C	C	C	
Brown rust	C	C	C	C	
Crown rust					C
Net blotch			C	C	
<i>Rhynchosporium</i>			C	C	
<i>Drechslera tritici-repentis</i> (see Note 1)	C	C			

C = Control

Note 1: Control of *Drechslera tritici-repentis* is supported by limited data only. Yield response may be obtained in the absence of visual disease.

Resistance Management

Jenton contains pyraclostrobin a member of the QoI cross-resistance group. Jenton should be used preventatively and should not be relied on for its curative potential.

When curative activity is needed apply QoI fungicides in mixture with a suitable dose of a curative fungicide from a different cross-resistance group. Jenton meets this requirement as it contains fenpropimorph. To ensure the effectiveness of the fenpropimorph component as a partner, Jenton should always be applied at the full recommended rate.

Use Jenton as part of an Integrated Crop Management (ICM) strategy incorporating other methods of control, including where appropriate other fungicides with a different mode of action. You must not apply more than two foliar applications of QoI containing products to any cereal crop.



There is a significant risk of widespread resistance occurring in *Septoria tritici* populations in the UK. Strains of barley powdery mildew resistant to QoI's are common in the UK. Failure to follow resistance management action may result in reduced levels of disease control.

3. Crops

Jenton is approved for use in winter wheat, winter barley, spring wheat, spring barley and oats.

Time of Application

Apply Jenton at the start of foliar disease attack. A maximum of two applications can be applied per crop. Applications can be made up to and including flowering just complete in winter and spring wheat and up to and including emergence of ear just complete in winter and spring barley and oats.

Rate of Application

Apply 2 litres Jenton in 200 litres of water per hectare.

4. Mixing and Spraying

Apply as a MEDIUM spray, as defined by BCPC.

Half fill the spray tank with clean water and start the agitation. SHAKE THE CONTAINER WELL before use and pour in the required amount of product. Rinse any empty containers thoroughly and add rinsings to the spray tank. Add the remainder of the water and continue agitation until spraying is completed.

When tank mixes are to be used, each product should be added separately to the spray tank.

The following does not form part of the product label under the Control of Pesticides Regulations 1986.

With many products there is a general risk of resistance developing to the active ingredients. For this reason a change in activity cannot be ruled out. It is generally impossible to predict with certainty how resistance may develop because there are so many crop- and use- connected ways of influencing this. We therefore have to exclude liability for damage or loss attributable to any such resistance that may develop. To help minimise any loss in activity the BASF recommended rate should in all events be adhered to.

Numerous, particularly regional or regionally attributable, factors can influence the activity of the product. Examples include weather and soil conditions, crop plant varieties, crop rotation, treatment times, application amounts, admixture with other products, appearance of organisms resistant to active ingredients and spraying techniques. Under particular conditions a change in activity or damage to plants cannot be ruled out. The manufacturer or supplier is therefore unable to accept any liability in such circumstances. All goods supplied by us are of high grade and we believe them to be suitable, but as we cannot exercise control over their mixing or use or the weather conditions during and after application, which may affect the performance of the material, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use; but nothing should be deemed to exclude or restrict any liability upon us which cannot be excluded or restricted under the provisions of the Unfair Contract Terms Act 1977 or any similar applicable law.





**Section 6 of the Health and Safety at Work Act
Additional Product Safety Information**

The product label provides information on a specific pesticidal use of the product; do not use otherwise, unless you have assessed any potential hazard involved, the safety measures required and that the particular use has "off-label" approval or is otherwise permitted under the Control of Pesticides Regulations.

The information on this label is based on the best available information including data from test results.

Safety Data Sheet

To access the Safety Data Sheet for this product scan the QR code or use the weblink below.



http://www.agricentre.basf.co.uk/go/jenton_sds

Alternatively, contact your supplier.

SPECIMEN

